arXiv:1209.4530v1 [gr-qc] 20 Sep 2012

On extreme Kerr-throats and zero temperature black-hole horizons.

MARTIN REIRIS.

MAX PLANCK INSTITUTE FÜR GRAVITATIONPHYSIK.
ALBERT EINSTEIN INSTITUT - GERMANY.
e-mail: martin@aei.mpg.de.

We show that there are no vacuum black holes horizons S over asymptotically flat maximal axisymmetric data sets saturating the universal inequality $A(S) \geq 8\pi|J(S)|$, where $A(S)$ and $J(S)$ are the area and angular momentum of S respectively. As equality is known to be reached (on globally different data sets) when and only when the intrinsic and extrinsic geometry of S is that of an extreme Kerr-throat sphere which has zero surface gravity or "temperature", our statement could be rephrased following this heuristic, as the non-existence of black holes of zero temperature (and topological origin) over asymptotically flat maximal slices in a vacuum space-time. We also show the rigidity of the extreme Kerr-throat data, which allows us to investigate the phenomenon of formation of extreme Kerr-throats along sequences of data sets. We prove our results for data sets in the smooth category, a hypothesis that is not entirely a technicality.

1 Introduction.

The Penrose singularity theorem ([10], Ch. 9.5) states that, if a trapped surface exists on a non-compact and complete Cauchy surface Σ, to be concrete in a vacuum space-time with asymptotically flat ends, then the space-time is null geodesically incomplete. The conclusion, which is believed to be an unequivocal signature for the emergence of black holes, is one of the few general results linking data sets and dynamical black hole formation. Let us concentrate for the moment in vacuum, maximal, asymptotically flat and axisymmetric data sets and suppose that the trapped surface is isotopic to a sphere at infinity over an asymptotically flat end as we depict in Figure 2. Then the boundary of the trapped region [3] is known to be a stable marginally outer trapped surface (MOTS), called an *apparent horizon*, which in turn can be interpreted as a *quasilocalization* of the event horizon. On the other hand, it was shown in [6] that for data sets of this type, any orientable surface S satisfies the universal inequality

$$A(S) \geq 8\pi|J(S)| \tag{1}$$

where $A(S)$ is the area of S and $J(S)$ its Komar angular momentum,

$$J(S) := \frac{1}{8\pi}\int_S K(\xi, \varsigma)\, dA \tag{2}$$

In this expression ξ is the rotational Killing field, K is the second fundamental form of the data set and ς is a unit normal to S (unique up to a sign). In particular the inequality (1) holds for the apparent horizon and because the angular momentum depends only on the homology class $[S]$ of S ($\nabla^a(K_{ab}\xi^b) = 0$) it is concluded that the area of the apparent horizon is always greater or equal than 8π times the angular momentum of the respective asymptotically flat end. Still, the inequality (1) is valid for axisymmetric stable minimal surfaces over axisymmetric maximal data sets regardless of any global information like completitud or asymptotic flatness [6] . Moreover it is proved that for stable minimal surfaces on maximal data sets, the equality is achieved when and only when the extrinsic and intrinsic geometry of S (inside the space-time) is that of an extreme Kerr-throat sphere, which, as defined in [6] is the geometry of a sphere at "infinity" on the asymptotically cylindrical end of an extreme Kerr solution. This article partly investigates which type of maximal data sets accept embedded surfaces saturating the inequality (1). As we will see the quest has deep theoretical implications. Before we continue and because it is relevant and central for the rest of the article and to give a level of accuracy to the present discussion, let us introduce the geometry of extreme Kerr-throats and extreme Kerr-throat spheres in detail. We resume the discussion immediately thereafter.

Consider the family of Kerr-solutions in the Boyer-Lindquist coordinates

$$\tag{3} \begin{aligned} \mathbf{g} = &-\left(\frac{\Delta - a^2\sin^2\theta}{\Sigma}\right) dt^2 - \frac{a\sin^2\theta(r^2+a^2-\Delta)}{\Sigma}(dtd\varphi + d\varphi dt) \\ &+ \left(\frac{(r^2+a^2)^2 - \Delta a^2\sin^2\theta}{\Sigma}\right)\sin^2\theta\ d\varphi^2 + \frac{\Sigma}{\Delta}\ dr^2 + \Sigma\ d\theta^2, \end{aligned}$$

where $\Sigma = r^2 + a^2\cos^2\theta,\ \Delta = r^2 + a^2 - 2mr$ and $a = J/m$ with m the mass. The Kerr *black holes* correspond to the range of parameters $m^2 \geq |J|$. The extreme Kerr-solutions corresponds to the case of equality, namely when $m^2 = |J|$ (or $a^2 = m^2$). Note that in this case $\Delta = (r-m)^2,\ r > m$. The extreme Kerr solutions have one asymptotically flat end (when $r \uparrow \infty$) and one cylindrical end (when $r \downarrow m$). All this is represented in Figure 1. The asymptotic cylinder has a well defined smooth data $(S^2\times\mathbb{R}, g_{\rm T}, K_{\rm T})$, indexed here by T from "Throat" and called the *extreme kerr-throat of angular momentum* $|J|$ (or of mass $m = \sqrt{|J|}$). It is given explicitly by

$$\tag{4} g_{\rm T} = \frac{4|J|\sin^2\theta}{1+\cos^2\theta}d\varphi^2 + |J|(1+\cos^2\theta)d\theta^2 + |J|(1+\cos^2\theta)d\tilde{r}^2$$

$$\tag{5} K_{\rm T} = \frac{2\sqrt{|J|}\sin^2\theta}{(1+\cos^2\theta)^{\frac{3}{2}}}(d\varphi d\tilde{r} + d\tilde{r}d\varphi)$$

The vector field $\partial_{\tilde{r}}$ is a Killing field and $|\partial_{\tilde{r}}|^2 = |J|(1+\cos^2\theta)$. Therefore $\partial_{\tilde{r}} = \alpha_{\rm T}\varsigma$ where $\alpha_T := \sqrt{|J|(1+\cos^2\theta)}$ and ς is a unit field normal to the spheres of constant $\tilde{r}$. The function α_T, which depends on the parametric factor $\sqrt{|J|}$, will play a fundamental role later on. The spheres of constant $\tilde{r}$ are called the *extreme Kerr-throat spheres* of area $A = 8\pi|J|$ (or of mass $m = \sqrt{A/8\pi}$). They are totally geodesic, i.e. the second fundamental form Θ is zero, and in particular minimal. Moreover they are stable minimal surfaces and the second variation of the area is zero when and only when it is in a direction proportional to $\partial_{\tilde{r}} = \alpha_T \varsigma$ (i.e. $\partial_{\tilde{r}}$ up to a factor). If we denote, as we will do later, with subindex $A, B, \ldots$ the coordinates $\{\theta, \varphi\}$ and with $i, j, \ldots$ the coordinates $\{\tilde{r}, \theta, \varphi\}$ then the data $\{g_{T,AB}, \Theta_{T,AB} = 0, K_{T,ij}\}$ will be called the *intrinsic and extrinsic geometry* of the extreme Kerr-throat sphere (of the given area). Note that the data of the extreme Kerr-throats (resp. extreme Kerr-throat spheres) are parametrized by their angular momentum $|J|$ (resp. area A) which plays the role of a global scale parameter.



Figure 1

Continuing with the discussion we observe from the universal inequality (1) that if an apparent horizon saturates the inequality (1), then, because $J(S)$ depends only on the homology class $[S]$ of S, it has to be a stable minimal surface. As said, there is uniqueness of such surfaces in the sense that in *areal coordinates* (see Section 2.3) the intrinsic and extrinsic geometry of S is that of an extreme Kerr-throat sphere of area $A(S) = 8\pi|J(S)|$. In this sense black hole horizons saturating the inequality (1) are unique. There is a worth mentioning interpretation for this type of black holes in terms of the well known thermodynamical heuristic from which we will borrow the terminology "*zero temperature black hole*". To explain, restrict attention to the family of Kerr solutions as seen over the maximal axisymmetric slice $\{t = 0\}$ in the Boyer-Lindquist coordinates. Non extremal Kerr solutions have positive temperature, given by

$$\frac{1 - (\frac{8\pi J}{A})^2}{(\frac{A}{4\pi} + \frac{16\pi J^2}{A})^{\frac{1}{2}}} \tag{6}$$

and a "localized" horizon, while the extremal solution has zero temperature and "non-localized" horizon, or, to phrase, an *asymptotic horizon* at the "end" of the cylindrical end. If for non-stationary or dynamical black holes, the temperature of the horizon is defined tentatively through (6) then, according to the mentioned uniqueness, zero temperature black hole horizons would have a unique intrinsic and extrinsic geometry, given by an extreme Kerr-throat sphere of a particular area. In this sense at least, zero temperature black hole horizons, of a given area, are unequivocally defined. We will keep using the terminology.

In the context of the discussion above, it is thus pertinent to ask whether zero temperature black hole horizons can arise on general axisymmetric and asymptotically flat maximal vacuum data sets or if on the contrary they can not, but could arise, as in the extreme Kerr solutions, only as asymptotic horizons on cylindrical ends. The present article investigates such situation and further related topics. Our main result in Theorem 1 shows that no black hole horizons of zero temperature exist, indeed no surface exists saturating (1), on axisymmetric and asymptotically flat, maximal vacuum data sets.

Theorem 1 *Let* $(\Sigma; g, K)$ *be a smooth vacuum axisymmetric maximal data set with possibly many asymptotically flat ends* $E_1, \ldots, E_n$*. Let* S *be any compact oriented surface. Then*

$$A(S) > 8\pi|J(S)| \tag{7}$$

where $A(S)$ *is the area of* S *and* $J(S)$ *its angular momentum.*

A representation of the type of data considered in the hypothesis of Theorem 1 is given in Figure 2. An immediate corollary is

Corollary 1 (Non-existence of zero-temperature black hole horizons) *There are no black hole apparent horizons, over smooth maximal axisymmetric and asymptotically flat vacuum data sets, of zero temperature.*

We will also prove a rigidity of extreme Kerr-throats, whose proof will follow basically the same lines as in Theorem 1.

Theorem 2 (Rigidity of extreme Kerr-throats) *Let* $(\Sigma; g, K)$ *be an homogeneously regular smooth axisymmetric maximal vacuum data set where* Σ *is diffeomorphic to* $S^2 \times \mathbb{R}$*. Suppose that for any sphere* S *isotopic to the factor* S^2 *we have* $A(S) \geq 8\pi|J|$ *where* $J = J(S) = J([S^2])$*, and suppose that there is at least one sphere* S_H *isotopic to the factor* S^2 *with* $A(S_H) = 8\pi|J|$*. Then* $(\Sigma; g, K)$ *is the extreme Kerr-throat data set (4)-(5) of angular momentum* $|J|$ *and* S_H *the extreme Kerr-throat sphere of area* $A(S_H) = 8\pi|J|$*.*

Theorem 2 is a remarkable manifestation, like in the positive mass theorem, of the nonlinearity of the constraint equations. The notion of *homogeneously regular* manifold which is explained in Section 2.4 essentially says that the metric is controlled in C^2 (on certain coordinates) on every metric ball of a uniform radius. We do not know at the moment if this condition can be removed and if only completitude of the data set is enough (it would be nice to answer this question). It seems however feasible to prove an optimal



Figure 2: Two possible configurations of topological black holes with the horizons marked with a S. On the left there is only one asymptotically flat end E_1, while on the right there are two, E_1 and E_2. Described are also large convex spheres on every asymptotically flat end. The horizon on the left has zero angular momentum because it encloses a compact boundary. The other surface shown on the left as well as the horizon shown on the right can have a priori non-zero angular momentum.

version of Theorem 2 prescinding of the topological condition $\Sigma \approx S^2 \times \mathbb{R}$. An interesting consequence of Theorem 2 is the following corollary on the formation of Kerr-throats.

Corollary 2 (Formation of extreme Kerr-throats) *Let* $\{(\Sigma; g_i, K_i)\}$ *be a sequence of smooth maximal vacuum axisymmetric and asymptotically flat initial data sets and let* $\{S_i\}$ *be a sequence of spheres* ($S_i \subset \Sigma_i$)*. Suppose that the sequence of data sets converges smoothly (and in the pointed sense) into a homogeneously regular maximal data set* $(\Sigma_\infty; g_\infty, K_\infty)$ *and that the sequence of spheres converges into an sphere* $S_\infty \subset \Sigma_\infty$*. If* Σ_∞ *is diffeomorphic to* $S^2 \times \mathbb{R}$ *and* $A(S_\infty) = 8\pi|J(S_\infty)|$*, then the limit data* $(\Sigma_\infty; g_\infty, K_\infty)$ *is the extreme Kerr-throat of angular momentum* $|J(S_\infty)|$ *and* S_∞ *is an extreme Kerr-throat sphere of area* $A(S_\infty)$*.*

Roughly speaking, under basic assumptions, a sequence of asymptotically flat, maximal axisymmetric data $\{(\Sigma_i; g_i, K_i)\}$ and sequence of embedded spheres $\{S_i\}$ can asymptotically saturate the universal inequality (1) only at the expense of the formation of an extreme Kerr-throat. More heuristically: extreme Kerr-throats form as the "temperature decreases to zero". A graphic realization of the phenomenon is given in Figure 3.

It is interesting to see Theorem 1 and Corollary 2 at work in the Kerr family of black holes. For the discussion that follows it is worth to keep in mind the Penrose diagram in Figure 1 of the maximal globally hyperbolic extension of the Kerr black holes for $0 < J^2 < m^4$. To facilitate the discussion we will take $m = 1$, therefore $0 < a = J^2 < 1$. The maximally globally hyperbolic space-time is represented in the Penrose diagram in four sectors. The quadrant on the right corresponds to the metric (3) for the range of coordinates

$$\{-\infty < t < \infty,\ 1 + (1 - a^2)^{\frac{1}{2}} < r < \infty,\ 0 < \theta < \pi,\ 0 < \varphi < 2\pi\}$$



Figure 3: The phenomenon of the formation of extreme Kerr-throats along sequence of data sets. The inequality (1) is saturated asymptotically along the spheres $\{S_i\}$.

and the upper quadrant corresponds to the range of coordinates

$$\{-\infty < t < \infty,\ 1-(1-a^2)^{\frac{1}{2}} < r < 1+(1-a^2)^{\frac{1}{2}},\ 0<\theta<\pi,\ 0<\varphi<2\pi\}$$

Note that in this case ∂_t is space-like while ∂_r is time-like. One can check directly Theorem 1 for the $\{t=0\}$ maximal slice which has the black hole horizon located at $\{r = 1+(1-a^2)^{\frac{1}{2}}\}$. The area is

$$A = (1+(1-a^2)^{\frac{1}{2}})8\pi$$

which is always greater than 8π and converges to 8π as $a \uparrow 1$. A more interesting phenomenon occurs when we see Corollary 2 in the light of the maximal slices given by the evolution in the maximal gauge of the initial slice. The maximal foliation penetrates inside the black hole region (the upper quadrant) and approaches in the long time limit to a maximal slice, "the last slice", lying entirely inside the black hole region[1] (See Figure 1). One observes now that the area of the spheres of constant r along the hypersurface $\{t = 1+(1-a^2)^{\frac{1}{2}}\}$ in the upper quadrant, evolve from $(1+(1-a^2)^{\frac{1}{2}})8\pi$ to $(1-(1-a^2)^{\frac{1}{2}})8\pi$. The area of the sphere formed by the intersection of the "last slice" and $\{t = 1+(1-a^2)^{\frac{1}{2}}\}$ thus approaches to 8π as $a \uparrow 1$. According to Corollary 2 the "last slice" must approach the Kerr-throat as $a \uparrow 1$. This can be seen explicitly by studying carefully the metric (3) as $a \uparrow 1$. What occurs is a remarkable phenomenon. As $a \uparrow 1$ the metric (3) in the upper quadrant of the Penrose diagram converges into the following metric

---

[1] We do not know an explicit expression for the last slice.

$$
\begin{aligned}
\mathbf{g} = &-(1+\cos^2\bar{\theta})d\bar{t}^2 + (1+\cos^2\bar{\theta})((\tan\bar{t})\bar{r}d\bar{t} + d\bar{r})^2 \\
&+ \frac{4\sin^2\bar{\theta}}{(1+\cos^2\bar{\theta})}(\bar{r}d\bar{t} - \bar{\varphi})^2 + (1+\cos^2\bar{\theta})^2 d\bar{\theta}^2
\end{aligned}
\tag{8}
$$

where we have barred the space-time coordinates to emphasize that they are not the same as the coordinates $\{t, r, \theta, \varphi\}$ which indeed degenerate as $a \uparrow 1$. This metric is a *metric soliton*[2] in the sense that, as seen as a flow $(g, K; N, X)(t)$ over $S^2 \times \mathbb{R}$ with coordinates $\{\bar{r}, \bar{\theta}, \bar{\varphi}\}$, we have

$$
\begin{aligned}
&g(\bar{t}) = g_T, \\
&K(\bar{t}) = K_T + (1+\cos^2\bar{\theta})^{\frac{1}{2}}(\tan\bar{t})d\bar{r}^2, \\
&N(\bar{t}) = (1+\cos^2\bar{\theta})^{\frac{1}{2}}, \\
&X(\bar{t}) = \bar{r}(\tan\bar{t})\partial_{\bar{r}} - \bar{r}\partial_{\bar{\varphi}}
\end{aligned}
$$

where, as it is apparent, the metric g does not evolve and remains equal to the three-metric g_T of the Kerr-throat given by (4). The only slice $\{\bar{t} = \mathit{const.}\}$ with zero mean curvature is $\{\bar{t} = 0\}$ and is the limit of the "last slices" as $a \uparrow 1$. As $\tan(\bar{t} = 0) = 0$ we conclude from the expression above that the data (g, K) over the slice $\{\bar{t} = 0\}$ is exactly the Kerr-throat. Note that $t \in (-\pi/2, \pi/2)$. The metric (8) is globally hyperbolic and partly coincides with the so called *near horizon geometry* (see [2], Eq. (2.5)) that has been extensively studied in the literature and is not globally hyperbolic. The spacetime described by the metric (8) has the remarkable property that is foliated by marginally trapped spheres saturating (1). In a certain sense the whole solution is a horizon.

To obtain the expression (8) make the change of variables $\{t, r, \theta, \varphi\} \to \{\bar{t}, \bar{r}, \bar{\theta}, \bar{\varphi}\}$

$$
\bar{t} = \frac{a(-\Delta)^{\frac{1}{2}}}{r^2 + a^2}t,\ \bar{r} = \arcsin\frac{1-r}{(1-a^2)^{\frac{1}{2}}},\ \bar{\theta} = \theta,\ \bar{\varphi} = \varphi - \frac{a}{r^2+a^2}t
$$

in the expression (3) and take the limit as $a \uparrow 1$.

We give now a glimpse of the basic idea behind the proof of Theorem 1. The idea is indeed rather simple. Suppose by contradiction that there is a surface $S = S_H$, under the hypothesis of Theorem 1, saturating (1). To be concrete we can think that S_H is an apparent horizon as discussed before. Then, as discussed before S_H has to be an extreme Kerr-throat sphere. Moreover as is proved in Proposition 3, unless the maximal lapse N_m (the solution of the Lapse equation (13) asymptotically one) is proportional[3] to α_T over S_H, then the area of the surface S_H strictly decreases along the evolution in the maximal gauge with zero shift. As $J(S_H)$ is preserved, then the universal inequality (1), which is valid for asymptotically flat data sets, would be violated in short times. Unfortunately this argument is based on the assumption that N_m is not proportional to α_T over S_H. It takes some technical work (and a large part of this article) to provide a proof in the spirit described but not relying on this singular assumption. The proof of Theorem 2 is also based on this simple idea. We will be explaining more along the article.

---

[2] This terminology is ours.

[3] As shown in Section 2.3, given any axisymmetric sphere S there are always areal-coordinates $\{\theta, \varphi\}$ which are unique up to the addition of a constant on φ. This implies that any continuous function $f : [0, \pi] \to \mathbb{R}$ can be thought as a continuous function $f(\theta)$ on S in a unique way.

It is rather remarkable that the non-existence of data sets with zero-temperature black hole horizons, ultimately a data set with certain properties, is ruled out using the dynamical Einstein equations. In this article we work in the smooth (C^∞) class of initial data and, at least in the approach of this article, its necessity is evident for instance in the use of the *unique continuation principle* [4] for elliptic equations recalled in Proposition 4. In the smooth class then, the dynamical equations (known to be well posed) give global information on data sets. However for low regularity, there is a chance to reverse the implications. Namely, to obtain information on the "minimal" regularity for which the dynamical equations in a particular gauge are well posed. This, a bit unlikely circumstance, is well illustrated in the following scenario. Suppose there exists a smooth maximal axisymmetric initial data $(\Sigma; g, K)$, $\Sigma \approx S^2 \times [0, \infty)$, and with boundary an extreme Kerr-throat sphere. Then regardless of the curvature at the boundary, the doubling of $(\Sigma; g, K)$ through its boundary, would be a data set in $H^2 \times H^1$. Now, if such a state exists then one could follow the same type of arguments leading to the proof of Theorem 1 to conclude that either,

1. The axisymmetric Einstein equations in the maximal gauge are not-well posed in $H^2 \times H^1$, or,
2. The inequality (1) does not hold on some asymptotically flat maximal data sets with $H^2 \times H^1$ regularity.

Motivated by these elaborations we state the following problem.

Problem 1 *Prove (or disprove) that there are no smooth maximal axisymmetric and asymptotically flat data sets* $(\Sigma; g, k)$ *with* Σ *diffeomorphic to* $S^2 \times [0, \infty)$ *and* $\partial\Sigma$ *an extreme Kerr-throat sphere.*

Note that by proving Problem 1 one would prove that there are no zero-temperature maximal axisymmetric black holes enclosing a material body, namely when the matter is strictly inside the black hole. We note finally that one can find maximal axisymmetric space-like sections Σ (with boundary) of the extended extreme Kerr solution diffeomoprhic to $S^2 \times [0, 1]$ where $S^2 \times \{0\}$ lies over the horizon, and therefore saturates the inequality (1), and where $S^2 \times \{1\}$ is a large convex sphere. However the induced state $(\Sigma; g, K)$ cannot be embedded into an axisymmetric and asymptotically flat maximal state and moreover the sphere $S^2 \times \{0\}$ is not totally geodesic in Σ.

The article is organized as follows. In Section 2 we recall the very basic notions and definitions required to read the article. Section 3 contains the proof of Theorems 1 and 2. In the Appendix we prove some propositions used in the proofs of the main results whose contents are a bit apart from the mainstream of the article.

2 Basic notions.

In this section we introduce the basic notions that we will use during the article. First we recall the Einstein vacuum equations from the dynamical point of view. This point of view is the appropriate one to analyze initial data as we will do in Theorems 1 and 2. Then we recall the second variation formula for minimal surfaces on three manifolds and the form that they acquire when the metric is the metric of an initial data for vacuum solutions. After that we introduce *areal* coordinates on axisymmetric spheres and other coordinates which play a crucial role in certain formulae. Finally we discuss basic but important properties of stable surfaces in axisymmetric three manifolds whose proofs are given in the Appendix. The presentation is somehow general and could be of use in further research.

2.1 The vacuum Einstein equations.

Let Σ be a fixed smooth manifold (say covered by a set of fixed charts $\{\{x^i, i = 1, 2, 3\}\}$). Given a Riemannian metric g we will denote by Ric and R the Ricci and the scalar curvatures of g respectively. Given a symmetric two tensor K, we will use the notation $|K|^2 = K_{ij}K^{ij}$ and $k = tr_g K = K_{ij}g^{ij}$. Finally given a vector field $X = X^i$, $\mathcal{L}_X$ will denote the Lie derivative along X.

Let $(g, K; N, X)(t)$ be a smooth flow on Σ of:

Riemannian metrics $g(t) = g_{ij}(t)$ and symmetric two-tensors $K(t) = K_{ij}(t)$,
Lapse functions $N(t) > 0$ and Shift vector fields $X(t) = X^i(t)$.

If for all t (in its range I) we have

$$\dot{g}_{ij} = -2NK_{ij} + (\mathcal{L}_X g)_{ij}, \tag{9}$$

$$\dot{K}_{ij} = -\nabla_i\nabla_j N + N(Ric_{ij} + kK_{ij} - 2K_i{}^k K_{kj}) + (\mathcal{L}_X K)_{ij}, \tag{10}$$

$$R = |K|^2 - k^2, \tag{11}$$

$$\nabla^i K_{ij} = \nabla_j k, \tag{12}$$

then the $3 + 1$ metric

$$\mathbf{g} = -(N^2 - X_i X^i)dt^2 + X_i(dt \otimes dx^i + dx^i \otimes dt) + g_{ij}dx^i dx^j$$

is a solution of the Einstein vacuum equations ($\mathbf{Ric} = 0$) on $I \times \Sigma$ where I is the interval on which t varies. If we let $\mathfrak{n}$ be a unit-normal to the level sets of the time coordinate (which are space-like hypersurfaces) we have $\partial_t = N\mathfrak{n} + X^i\partial_i$. Moreover $K(t)$ are the second fundamental forms of the level sets of the time coordinate, namely of $\{t\} \times \Sigma$. Equations (9)-(10) are the *dynamical equations* and (11)-(12) are the *energy* and *momentum constraints equations* respectively. The evolution is said to be *maximal* if $k(t) = 0$ for all t. In such case the Lapse satisfies, at any time, the *Lapse equation*

$$\Delta N = |K|^2 N \tag{13}$$

Conversely if Σ is a space-like manifold (possibly with boundary and covered by a set of coordinate charts $\{\{x^i, i=1,2,3\}\}$) on a vacuum space-time and V is a non-zero time-like vector field defined on an open space-time neighborhood of Σ then one can obtain a flow $\Sigma(t)$, at least for a short time, by moving Σ along V. Coordinates charts $\{x^i, i=1,2,3\}$ are propagated by V to every $\Sigma(t)$ and any two $\Sigma(t)$ and $\Sigma(t')$ are canonically diffeomorphic. In this way one obtains a flow $(g_{ij}(t), K_{ij}(t))$ for the induced three-metrics and second fundamental forms on the *fixed* manifold Σ. Writing $V|_{\Sigma(t)} = N(t)\mathfrak{n} + X^i(t)\partial_i$, where $\mathfrak{n}$ is a space-time unit normal to $\Sigma(t)$, then one obtains a flow of Lapse functions $N(t)$ and Shift vectors $X(t) = X^i\partial_i$. The flow $(g(t), K(t); N(t), X(t))$ satisfies (9)-(12). *It is important to stress finally that the flow is on a fixed manifold* Σ.

2.2 Minimal surfaces and the second variation of area.

Let $(\Sigma; g, K)$ be a data set. An embedded surface S is minimal if its mean curvature is identically zero. Suppose that S is orientable and minimal. Let ς be a unit normal vector field to S and let $\alpha : S \to \mathbb{R}$ be a smooth function. Then, the second variation of the area, A''_α, for the deformation of S along $\alpha\varsigma$ is the well known [5]

$$A''_\alpha(S) = \int_S |d\alpha|^2 - (|\Theta|^2 + Ric(\varsigma,\varsigma))\alpha^2 dA \tag{14}$$

where Θ is the second fundamental form of S. The mean curvature will be denoted by $\mathrm{tr}_h\Theta$. The minimal surface S is said to be *stable* if $A''_\alpha(S) \geq 0$ for all α. In dimension three we further have the identity $2\mathcal{K} = (\mathrm{tr}_h\Theta)^2 - |\Theta|^2 + R - 2Ric(\varsigma,\varsigma)$, where $\mathcal{K}$ is the Gaussian curvature of S and when S is minimal $\mathrm{tr}_h\Theta = 0$. From this and the energy constraint we obtain

$$\int_S (|d\alpha|^2 + \mathcal{K}\alpha^2)\ dA \geq \frac{1}{2}\int_S (|\Theta|^2 + |K|^2 - k^2)\alpha^2\ dA \tag{15}$$

2.3 Areal and polar coordinates for axisymmetric spheres and further coordinates.

Let (Σ, g) be an a axisymmetric manifold. Then the group $U(1)$ acts on (Σ, g) and the orbits are either circles or fixed points. The set of fixed points are a set of complete one dimensional manifolds (i.e. diffeomorphic to lines or circles) which we call the *axes* and denote by $\{\mathcal{P}\}$.

I. *Areal coordinates on spheres.* Let $S \subset \Sigma$ be an axisymmetric sphere. Then S is foliated by $U(1)$-orbits $\{C\}$. Two of them are just points (the poles), and denote them (arbitrarily) by $\mathcal{N}$ (from "North") and $\mathcal{S}$ (from "South"). Every orbit which is not a pole divides S into two discs, denote them by $D^{\mathcal{N}}(C)$ and $D^{\mathcal{S}}(C)$ (the first contains $\mathcal{N}$ and the second $\mathcal{S}$). Denote $A^{\mathcal{N}}(C) = \text{Area}(D^{\mathcal{N}}(C))$ and $A^{\mathcal{S}}(C) = \text{Area}(D^{\mathcal{S}}(C))$. Define the polar coordinate $\theta(C)$ at C by

$$A^{\mathcal{N}}(C) = \frac{A(S)}{2}\big(1 - \cos\theta(C)\big)$$

To define an azimuthal coordinate φ proceed as follows. The gradient of θ (inside S) defines a vector field perpendicular to the orbits and invariant under the $U(1)$-action. The integral curves foliate smoothly $S \setminus \{\mathcal{N}, \mathcal{S}\}$. Denote the foliation by $\{C'\}$. Pick any integral curve to define $\{\varphi = 0\}$. Then $\varphi(C')$ is the angle necessary to rotate $\{\varphi = 0\}$ to get C'. In these coordinates the induced metric h (from g) on S looks like

$$h = \left(\frac{A(S)}{4\pi}\right)^2 e^{-\sigma} d\theta^2 + e^{\sigma} \sin^2\theta d\varphi^2 \tag{16}$$

where $\sigma = \sigma(\theta)$. Note that $dA = (A(S)/4\pi)\sin\theta d\theta d\varphi$.

II. *Polar coordinates on discs.* In addition to areal coordinates we will use polar coordinates for discs $D^{\mathcal{N}}$ or $D^{\mathcal{S}}$. Around the respective pole the induced metric h in polar coordinates is

$$h = ds^2 + \left(\frac{\ell(s)}{2\pi}\right)^2 d\varphi^2 \tag{17}$$

Of course $\{s = 0\}$ is the pole and $\ell(s)$ is the length of the orbit $C(s)$ at a h-distance s from the pole. Say $\{s = 0\}$ is the north pole. Then we will denote $A(s) := A^{\mathcal{N}}(C(s))$. This notation will be used later (see the proof of Proposition 5). In particular $A(s) = \int_0^s \ell(\bar{s})d\bar{s}$.

III. *Gaussian-coordinates around surfaces.* Let S be an axisymmetric sphere provided with areal-coordinates $\{\theta, \varphi\}$. For $\bar{r} > 0$ small enough let

$$\mathcal{T}(S, \bar{r}) := \{p \in \Sigma, dist(p, S) \leq \bar{r}\}$$

be the tubular (closed) neighborhood of S of radius $\bar{r}$. For every $q \in S$ let $\gamma_q(r)$ be the g-geodesic emanating from q, perpendicular to S and parametrized with (signed) arc-length $r \in [-\bar{r}, \bar{r}]$. For every $p \in \mathcal{T}(S, \bar{r})$ let $q(p)$ be the initial point in S such that $\gamma_{q(p)}(r(p)) = p$ ($|r(p)| = dist_g(p, S)$). The g-*Gaussian coordinates* on $\mathcal{T}(S, \bar{r})$, $\{r, \theta, \varphi\}$, are defined through

$$(r, \theta, \varphi)(p) = (r(p), \theta(q(p)), \varphi(q(p)))$$

We will use Gaussian coordinates often in this article. Observe that ∂_φ is the rotational Killing field. In Gaussian coordinates the metric g is written as

$$g = dr^2 + h$$

where $h(\partial_r, -) = h(-, \partial_r) = 0$ at any point.

IV. *Space-time coordinates.* The Gaussian coordinates are propagated along the evolution (using (N, X)) in such a way that $\{(r, \theta, \varphi, t)\}$ (t small) is a space-time coordinate system around S. We will use the index $A, B, \ldots$ when we use only the two coordinates (θ, φ) while we will use the index $i, j, \ldots$ when use the three coordinates (r, θ, φ). In this sense the components of h are h_{AB} while those of g are g_{ij}.

Above we introduced $\mathcal{T}(S, \bar{r})$. More in general, we will use

$$\mathcal{T}(\Omega, \bar{r}) = \{p \in \Sigma / \mathrm{dist}_g(p, \Omega) \leq \bar{r}\}$$

to denote the (closed) tubular neighborhood of a set $\Omega \subset \Sigma$ and radius $\bar{r}$. We will also write $\mathcal{T}_g(\Omega, \bar{r})$ whenever it is necessary to stress that the tubular neighborhood is with respect to the particular metric g.

2.4 Homogeneously regular manifolds.

A complete manifold is homogeneously regular if the injectivity radius is uniformly bounded from below (away from zero) and the curvature is uniformly bounded from above. A more quantitative but equivalent definition is the following [8].

A complete manifold Σ is ρ_0-homogeneously regular if there is $0 < \nu < 1/2$ such that at any point $p \in \Sigma$ we have

1. *the exponential map Exp, from $B_{T_p\Sigma}(p, \rho_0)$ into $B(p, \rho_0)$ is a diffeomorphism (where $B_{T_p\Sigma}(p, \rho_0)$ is the ball in the tangent space $T_p\Sigma$ of center p and radius ρ_0 and $B(p, \rho_0)$ is the geodesic ball in Σ of center p and radius ρ_0).*
2. *Let $\{(\bar{x}^1, \bar{x}^2, \bar{x}^3)\}$ be cartesian coordinates in $T_p\Sigma$. Define as usual coordinates on $B(p, \rho_0)$ by $(x^1, x^2, x^3) = (\bar{x}^1, \bar{x}^2, \bar{x}^3) \circ Exp^{-1}$. Then,*

$$|g_{ij} - \delta_{ij}| \leq \nu, \quad \sup_{B(p,\rho_0)} |\frac{\partial g_{ij}}{\partial x^k}| \leq \frac{\nu}{\rho_0}, \quad \sup_{B(p,\rho_0)} |\frac{\partial g_{ij}}{\partial x^k x^l}| \leq \frac{\nu}{\rho_0^2}$$

for all i, j, k, l in $\{1, 2, 3\}$.

Essentially, a homogeneously regular manifold is one for which the metric is controlled in C^2 (on certain coordinates) on every metric ball of a uniform radius. It is easy to see from the definition that if (Σ, g) is ρ_0-homogeneously regular then $(\Sigma, \lambda_0^2 g)$, $\lambda_0 > 0$, is $\lambda_0\rho_0$-homogeneously regular. In particular, for any $\epsilon > 0$ there is $\lambda_0(\epsilon, \rho_0)$ such that for any point p, the scaled metric $\lambda_0^2 g$ on $B_{\lambda_0^2 g}(p, 1)$ is ϵ-close in C^2 (in the coordinates described above) to the flat metric on the unit ball of $\mathbb{R}^3$. Before $B_{\lambda_0^2 g}(p, 1)$ is the ball of center p and radius one with respect to $\lambda_0^2 g$.

For a manifold with boundary the definition is similar, but we require that there is an extension $(\bar{\Sigma}, \bar{g})$ of (Σ, g) such that the boundary of $\bar{\Sigma}$ lies at a $\bar{g}$-distance greater than ρ_0 from $\partial\Sigma$ and at every point p of Σ the points 1 and 2 hold. Of course every compact manifold, with boundary or not, is ρ_0-homogeneously regular for some ρ_0.

2.5 Minimal surfaces in axisymmetric spaces.

Let $B(o,r)$ be the ball in $\mathbb{R}^3$ of radius r and centered at the origin $\{o=(0,0,0)\}$. We will think $\mathbb{R}^3$ as an axisymmetric space-time where the $U(1)$-action is by rotations around the z-axis. Every smooth connected axisymmetric surface S, possibly with boundary, inside $\overline{B(o,1)}$ (with boundary not necessarily in $\partial B(o,1)$) is either

$\mathcal{I}_0$. an axisymmetric sphere or an axisymmetric torus (zero boundary component),

$\mathcal{I}_1$. an axisymmetric disc (one boundary component),

$\mathcal{I}_2$. an axisymmetric cylinder, namely diffeomorphic to $S^1\times[0,1]$ (two boundary components),

Let $\mathscr{S}_i$, $i=0,1,2$, be the set of connected axisymmetric surfaces embedded in $B(o,1)$ of type $\mathcal{I}_i,\ i=0,1,2$ respectively and with boundary, if any, lying in $\partial B(o,1)$. Let C be an orbit in $B(o,1)$ which is not a point in the axis. Let $\mathscr{S}_1^C$ be the set of axisymmetric discs in $\mathcal{I}_1$ with boundary C.

The following Proposition is straightforward (indeed the first *item* is trivial) from standard properties of minimal surfaces. As it is standard and the proof has few to do with the content of the article we divert the proof until the Appendix.

Proposition 1 *There is* $0<L<1$ *such that*

1. *For every orbit* $C\subset B(o,L)$*, which is not a point in the axis, there is a unique disc* D *in* $\mathscr{S}_1^C$ *which is minimal. Such unique disc minimizes area among all the surfaces in the family of surfaces* $\mathscr{S}_1^C$ *and therefore is stable.*
2. *There are no stable axisymmetric minimal surfaces in the family of surfaces* $\mathscr{S}_0\cup\mathscr{S}_2$ *and intersecting* $B(o,L)$*.*

Of course, by scaling the catenoid, one can easily show that there are complete axisymmetric minimal surfaces in $B(o,1)$ reaching as close to the origin $\{o\}$ as wished, but they are not stable.

It will be necessary to dispose of a version of Proposition 1 but for C^2-perturbed axisymmetric metrics. More precisely

Proposition 2 *Let* g *be a smooth axisymmetric metric on* $B(o,2)\subset\mathbb{R}^3$*. Assume that the polar coordinate system* $\{(z,\rho,\varphi)\}$ *of* $\mathbb{R}^3$ *is adapted to* g *in the sense that, in these coordinates* g_{ij} *is independent on* φ*. Then there is* $\epsilon_0>0$ *and* $0<L<1$ *such that if* g *is* ϵ_0*-close in* C^2 *to the flat metric in* $\mathbb{R}^3$ *the following is true.*

1. *For every orbit* $C\subset B_g(o,L)$*, which is not a point in the axis, there is a unique disc* D *in* $\mathscr{S}_1^C$ *which is minimal. Such unique disc minimizes area among all the surfaces in the family* $\mathscr{S}_1^C$*.*
2. *There are no stable axisymmetric minimal surfaces of the family* $\mathscr{S}_0\cup\mathscr{S}_2$ *and intersecting* $B_g(o,L)$*.*

A corollary which we also prove in the Appendix is the following.

Corollary 3 *Let (Σ, g) be a ρ_0-homogeneously regular, non-necessarily compact axisymmetric three-manifold with smooth, compact and axisymmetric boundary. Assume that the (outward) mean curvature of the boundary is bounded below by $\mu_0 > 0$ and the norm of the second fundamental form bounded above by μ_1. Then, for any embedded stable axisymmetric minimal surface S in Σ we have*

1. $S \subset (\Sigma \setminus \mathcal{T}(\partial\Sigma, \epsilon_1(\rho_0, \mu_0, \mu_1)))$*, i.e. every S lies at a distance greater than ϵ_1 from $\partial\Sigma$.*
2. $A(S) \geq A_0(\rho_0, \mu_0, \mu_1) > 0$*, i.e. there is a uniform lower bound for the areas of the surfaces S.*
3. *There is $\epsilon_2(\rho_0, \mu_0, \mu_1) < \epsilon_1/2$ such that $\partial\big(\mathcal{T}(\{\mathcal{P}\}, \epsilon_2) \setminus \mathcal{T}(\partial\Sigma, \epsilon_1)\big)$ is smooth and foliated by orbits and for any one such orbit there is a unique area minimizing disc with boundary the orbit. Moreover, if S intersects $\mathcal{T}(\{\mathcal{P}\}, \epsilon_2)$ then it does at two of such discs. Thus, either S intersects the axes $\{\mathcal{P}\}$, in which case it does twice and the surface is an sphere, or it lies at a distance from $\{\mathcal{P}\}$ greater than ϵ_2.*
4. *Consider the adapted coordinates of any disc D as in item 3, namely where the metric is $h = ds^2 + (\ell/2\pi)^2(s)\ d\varphi^2$. Then s ranges in an interval $(0, s_D]$ and $s_D \in [s_0(\rho_0, \mu_0, \mu_1), s_1(\rho_0, \mu_0, \mu_1)]$. Moreover we have $|\ell(s) - 2\pi s| \leq c_0(\rho_0)s^2$.*

When Σ, ρ_0-homogeneously regular and complete but boundary-less then the items 2,3 and 4 hold but now the estimates depend only on ρ_0.

To finish this section let us mention that stable minimal surfaces in either asymptotically flat axisymmetric manifolds or in compact axisymmetric manifolds with axisymmetric boundary are necessarily axisymmetric. In particular one can replace the hypothesis "stable axisymmetric minimal surface" in Corollary 3 by just "stable minimal surface". We explain this known but important fact in what follows. Let Σ be a three-manifold of one of the two mentioned types and let S be a compact embedded stable minimal surface. Let again ξ denote the axial Killing field and ς a unit normal to S. Let $\lambda^N =< \xi, \varsigma >$ denote the normal component of ξ over S. If λ^N is nowhere zero then the $U(1)$-orbits are transversal to S at any of its points which implies that Σ must be diffeomorphic to $S \times S^1$. This is an impossibility. It follows that λ^N is somewhere zero. We show now that λ^N is identically zero and therefore invariant under the $U(1)$ action (thus axisymmetric). Suppose that λ^N is non-zero somewhere in S. As ξ is Killing we deduce that the second variation of the area along $\lambda^N\varsigma$ is zero. On the other hand as S is stable we deduce that the first eigenvalue of the the second variation operator[4] is zero and λ^N is the eigenfunction. But the eigenfunction of the first eigenvalue is always nowhere zero and we reach a contradiction.

---

[4] The second variation operator is here that obtained from varying (14) that is $L\phi = -\Delta\phi - (|\Theta|^2 + Ric(\varsigma, \varsigma))\phi$.

3 Proof of the main results.

3.1 The second variation of area in time for extreme Kerr-throat spheres.

The proofs of Theorems 1 and 2 rest on the crucial negativity of the second variation of the area in time of stable extreme Kerr-throat spheres (on maximal, axisymmetric initial data) when $N(0)$ is not proportional to α_T over the sphere and $X(0) = 0$. Due to the basic importance of the computation we present it below before going into the proofs of the main results.

Proposition 3 *Let* $(\Sigma; g, K)$ *be a vacuum, axisymmetric, maximal initial data set, Let* S_H *be a stable extreme Kerr-throat sphere. Suppose that the data evolves following the vacuum Einstein equations with smooth lapse and shift* $(N(t), X(t))$ *such that* $X(0) = 0$. *Then, at time zero we have*

$$\dot{A}(S_H) = 0, \tag{18}$$

$$\ddot{A}(S_H) = -A''_N(S_H) \leq 0. \tag{19}$$

Moreover, equality in the second inequality in (19) holds iff in areal-coordinates we have $N\big|_{S_H} = c\alpha_T$ *for a constant* c.

Note 1 *In a vacuum axisymmetric maximal and asymptotically flat data set, every extreme Kerr-throat sphere is necessarily stable. In Proposition 3 there is no global assumption on the initial data of any kind and the stability of* S_H *has to be imposed a priori. Nevertheless, as can be seen from the proof, even if* S_H *is not stable we still have* $\ddot{A}(S_H) = -A''(S_H)$.

Proof: In the following calculation one can use for instance the space-time coordinate system introduced in Section 2.3. We have

$$\dot{A}(S_H) = \frac{1}{2}\int_{S_H} \dot{h}_{AB}h^{AB}dA = \int_{S_H} -NK_{AB}h^{AB} + \frac{1}{2}(\nabla_A X_B + \nabla_B X_A)h^{AB}dA.$$

Recalling that at time zero we have $K_{AB}(0) = 0$, and $X(0) = 0$ then, at time zero we have

$$\ddot{A}(S_H) = \int_{S_H} -N\dot{K}_{AB}h^{AB} + \frac{1}{2}(\nabla_A \dot{X}_B + \nabla_B \dot{X}_A)h^{AB}dA.$$

As S_H is totally geodesic at time zero, we have

$$\int_{S_H} \frac{1}{2}(\nabla_A \dot{X}_B + \nabla_B \dot{X}_A)h^{AB}dA = \int_{S_H} Div_{S_H}\Pi(\dot{X})dA = 0.$$

where $\Pi(\dot{X})$ is the projection of $\dot{X}$ onto the tangent space of S_H, and Div_{S_H} is its divergence as a vector field on (S_H, h). Therefore, recalling that the initial data is maximal, that is $k = 0$, we have

$$\ddot{A}(S_H) = \int_{S_H} -N(-\nabla_A \nabla_B N + N(Ric_{AB} - 2K_{Ai}K^i{}_B))h^{AB} dA.$$

Again, S_H is totally geodesic at time zero, therefore

$$\int_{S_H} N(\nabla_A \nabla_B N) h^{AB} dA = -\int_{S_H} |dN|^2 dA.$$

where d, as before, is the differential of N in S_H.

On the other hand, $Ric_{AB}h^{AB} = R - Ric(\varsigma, \varsigma)$ and, because of the energy constraint and because that the only nonzero components of K_{ij} are $K_{\theta r}$ (and $K_{r\theta}$) then $2K_{Ai}K^i{}_B h^{AB} = |K|^2 = R$. Altogether we obtain, at time zero,

$$\ddot{A}(S_H) = \int_{S_H} -|dN|^2 + Ric(\varsigma, \varsigma)N^2 dA = -A''_N(S_H)$$

because the second fundamental form of S_H is zero.

Finally if S_H is stable then for any $\alpha : S_H \to \mathbb{R}$ we have $A''_{\alpha\varsigma}(S_H) \geq 0$. On the other hand we obtain equality in (15) if $\alpha = \alpha_T$ which implies that the right hand side of (14) is zero [6] and thus $A''_{\alpha_T \varsigma}(S_H) = 0$. From both facts we deduce that zero is the first eigenvalue of the stability operator with eigenfunction α_T. Moreover from the uniqueness of the first eigenfunction we have $A''_{\alpha\varsigma}(S_H) = 0$ iff α is proportional to α_T.

□

3.2 Proof of Theorem 1.

Let us recall the main argument behind the proof of Theorem 1. Assume by contradiction the existence of $(\Sigma; g_0, K_0)$ containing an extreme Kerr-throat sphere S_H and to fix ideas suppose that S_H is isotopic to a sphere at “infinity” on one of the asymptotically flat ends. As was discussed in the introduction this is the case for instance if S_H is an apparent horizon. In any case S_H divides Σ in two connected components. Suppose that the *maximal lapse* N_m, namely the solution to (13) which is asymptotically one at infinity on Σ, is not proportional to α_T over S_H. Then, evolving the initial data (g_0, K_0) in the maximal gauge with zero shift and using Proposition 3 and the conservation of angular momentum one obtains, in short time, a sphere on an asymptotically flat, axisymmetric, maximal data violating (1) which is impossible. Unfortunately it could be (a priori) that N_m is proportional to α_T over S_H and therefore the argument, as such, is incomplete. However one can technically modify it, still following the same ground idea, to make it work. The modification consists in working on a large but compact region of the initial data enclosed by large convex spheres, where as we will see using Proposition 4 it is always possible to find a positive solution N_0 of the Lapse equation not proportional to α_T over S_H. Then flow the initial data over such compact region along certain (see later) axisymmetric lapse and shift $(N, X)(t)$ with $N|_{t=0} = N_0$ and $X|_{t=0} = 0$. Unfortunately the flow will be known to be maximal only at time zero and at later times maximality could fail. This is an important drawback because although by Proposition 3 we have, for t small,

$$A(S_H) \leq 8\pi|J(S_H)| - \frac{A''_{N_0}(S_H)}{4}t^2, \quad A''_{N_0}(S_H) > 0 \tag{20}$$

(where here and below $A(S_H) := A_{g(t)}(S_H)$), the inequality $A(S) \geq 8\pi|J(S)|$ is not known to hold on non-maximal slices, and the original contradiction argument may be inapplicable. Here is where we use that the initial slice is maximal and that, because $N|_{t=0} = N_0$ satisfies the Lapse equation and $\dot{k}|_{t=0} = -\Delta N_0 + |K_0|^2 N_0 = 0$, then the mean curvature $k(t)$ at small times behaves as $k(t) \approx O(t^2)$. This order of failure of maximality allows us to prove in Proposition 5, and for small times, the lower estimate $A(S_t) \geq 8\pi|J(S_t)| - \Lambda_0 t^4$ for the area of stable axisymmetric minimal surfaces S_t. This is then used in the *Proof of Theorem 1* to obtain the inequality

$$A(S_H) \geq 8\pi|J(S_H)| + O(t^4) \tag{21}$$

for small times. Inequalities (20)-(21) show a contradiction in the original spirit. We move then to prove the preliminary Propositions 4 and 5, the proof of Theorem 1 is given afterwards.

Proposition 4 *Let $(\Omega; g, K)$ be a vacuum maximal state where Ω is a compact manifold with smooth boundary. Suppose that either*

A1. *There is an extreme Kerr-throat sphere S_H dividing Ω into two connected components Ω_1 and Ω_2, or,*

A2. *There are extreme Kerr-throat spheres S^1_H and S^2_H, the union of which divides Ω into two connected components Ω_1 and Ω_2.*

Then there is an axisymmetric solution N of the Lapse equation which is positive on $\Omega^\circ = \Omega \setminus \partial\Omega$ and

1. *is not proportional to α^1_T over S_H in case* **A1**, *or,*
2. *is not proportional to α^1_T over at least one of the surfaces S^1_H or S^2_H in case* **A2**.

where $\alpha^1_T := \sqrt{1+\cos^2\theta}$ in the areal-coordinates of the respective sphere.

Note 2 *If S_H is an extreme Kerr-throat sphere of area A_H then the function α_T, as defined in the introduction is, $\alpha_T = \sqrt{A_H/8\pi}\sqrt{1+\cos^2\theta}$. Therefore if N is not proportional to α^1_T over S_H then it not proportional either to α_T.*

Proof. Denote by W_1 and W_2 the set of connected components of $\partial\Omega$ belonging to Ω_1 and Ω_2 respectively. As Ω has non-empty boundary by assumption then W_1 and W_2 cannot be empty at the same time. We will assume that $W_2 \neq \emptyset$. We discuss cases **A1** and **A2** separately.

Case **A1**. Let $\bar{N}_a$ and $\bar{N}_b$ be two positive and linearly independent axisymmetric functions over W_2 and let N_a and N_b be, respectively, the solutions of the Lapse equation with boundary data

$$\begin{aligned}
&\Delta N = RN,\\
&N\big|_{W_2} = \bar{N}_a \text{ or } \bar{N}_b,\\
&N\big|_{W_1} = 0, \text{ if } W_1 \neq \emptyset
\end{aligned}$$

The uniqueness of the solutions N_a, N_b and the axisymmetry of the boundary data imply that N_a and N_b must coincide with their rotational averages and therefore must be rotational symmetric. If either N_a or N_b are not proportional to α^1_T over S we are done. On the other hand, if both N_a and N_b are proportional to α^1_T over S_H, we can consider a non-zero linear combination $c_a N_a + c_b N_b$ which is equal to zero over S_H. The combination is also zero over W_1 if $W_1 \neq \emptyset$ and as $\partial\Omega_1 = W_1 \cup S_H$ uniqueness implies that the combination is zero all over Ω_1. But because $\bar{N}_a$ and $\bar{N}_b$ are linearly independent then $c_a N_a + c_b N_b$ is not identically zero over Ω_2. This contradicts the *unique continuation principle for elliptic equations* [4]. Thus either N_a and N_b are not proportional to α^1_T over S_H.

Case **A2**. Let $\bar{N}_a$, $\bar{N}_b$ and $\bar{N}_c$ be three positive and linearly independent axisymmetric functions over W_2. Let N_a, N_b and N_c be the axisymmetric solutions to the Lapse equation with boundary data zero over W_1 if non-empty and boundary data $\bar{N}_a$, $\bar{N}_b$ or $\bar{N}_c$ over W_2, respectively. If all of N_a, N_b and N_c are proportional to α^1_T over both S^1_H and S^2_H then, again, one can easily find a linear combination of the three that is zero over both S^1_H and S^2_H. A contradiction is then found as in *Case* **A1**. □

Let $(\Sigma; g, k)$ be, as in Theorem 1, a maximal (vacuum and axisymmetric) data set with possibly many asymptotically flat ends $E_1, \ldots, E_n$, $n \geq 1$. Let $\mathfrak{S}_1, \ldots, \mathfrak{S}_n$ be large axisymmetric and strictly convex spheres on each of the ends $E_1, \ldots, E_n$ respectively. Suppose there is an extreme Kerr-throat sphere S_H. If S_H does not divide Σ into two connected components then one can cut Σ along S_H to get a manifold with two boundary components (say S^1_H and S^2_H) and glue back smoothly a copy of it (crossing the boundaries). The result is a smooth manifold with $2n$ asymptotically flat ends and having two embedded extreme Kerr-throat spheres S^1_H and S^2_H the union of which divides the manifold into two connected components. The conclusion is that if there is an extreme Kerr-throat sphere then either we are in the hypothesis **A1** of Proposition 4 or that we can construct a data set in the hypothesis **A2**. In any case we are denoting by Ω the regions enclosed by the large spheres (including the large spheres). We will continue as if we were either in the hypothesis **A1** or **A2** therefore.

We note now that there is always a positive solution N_0 to the Lapse equation on Ω which is not proportional to α_T over S_H (in case **A1**) or is not proportional to α_T over one of the spheres S^1_H or S^2_H (in case **A2**). Indeed let N_m be the (positive) solution to the Lapse equation that is asymptotically one over any asymptotically flat end. If N_m is proportional to α_T over S_H (in case **A1**) or is proportional to α_T over both, S^1_H and S^2_H (in case **A2**) then we can add to it a solution to the Lapse equation as in Proposition 4 to obtain a positive solution on Ω with the desired property. From now on let N_0 be the positive axisymmetric solution of (13) in Ω, that is not proportional to α_T over S_H (in case **A1**) or is not proportional to α_T over either S_H^1 or S_H^2 (in case **A2**).

Let $\mathfrak{n}$ be a (one of the two possible) unit-normal fields to Ω inside the space-time. For any $p \in \Omega$ let $\gamma_p(\tau)$ be the (space-time) geodesic emanating from p, in the direction of $\mathfrak{n}(p)$, and parametrized with arc length τ. In the domain $\{o/o = \gamma_p(\tau), p \in \Omega, 0 \leq \tau \leq \tau_0\}$, τ_0 a small constant, we consider a vector field V by

$$V(o) := N_0(p)\gamma_p'(\tau), \quad \text{if } \gamma_p(\tau) = o. \tag{22}$$

Flowing the domain Ω inside the space-time (generated by the data) by the the vector field V, we obtain an evolution flow $(g, K; N, X)(t)$ over the fixed Ω where t is the parameter associated to the flow by V (see Section 2.1). Moreover we have $X_{t=0} = 0$ and $N_{t=0} = N_0$. Note that the evolution induced by V is axisymmetric, namely $(g, K; N, X)$ are axisymmetric. However the evolution does not have to be maximal, that is, we do not necessarily have $k(t) = 0$ for all t. Despite of this we have

$$k\big|_{t=0} = 0, \quad \dot{k}\big|_{t=0} = -\Delta N_0 + RN_0 = 0.$$

Thus there is $k_0 > 0$ such that $|k(t)| \leq k_0 t^2$. Therefore the energy constraint implies (at time t) $R = |K|^2 + O(t^4)$. The following auxiliary proposition gives a crucial lower bound on the area $A(S_t)$ of stable minimal surfaces S_t in $(\Omega; g(t), K(t))$ (with an upper bound on their areas).

Proposition 5 *For any $A_1 > 0$ there are $t_0 > 0$ and $\Lambda_0 > 0$ such for any $0 < t < t_0$ and orientable stable axisymmetric minimal surface S_t on $(\Omega; g(t), K(t))$ with $A(S_t) \leq A_1$ we have*

$$A(S_t) \geq 8\pi|J(S_t)| - \Lambda_0 t^4$$

Proof. Let $\bar{t}_0$, ρ_0, μ_0 and μ_1 be such that for any t in $[0, \bar{t}_0]$ the manifold $(\Omega, g(t))$ is ρ_0-homogeneously regular and with strictly convex boundary of (outward) mean curvature greater or equal than μ_0 and norm of the second fundamental form bounded by μ_1. Let ϵ_2 and A_0 be as in Corollary 3. Let $k_0 > 0$ be (as before) such that for all $t \in [0, \bar{t}_0]$, we have $|k(t)| \leq k_0 t^2$. Finally let $\ell_0 = \sup\{2\pi|\xi(p)|_{g(t)}, p \in \Omega, t \in [0, \bar{t}_0]\}$.

For the proof we distinguish two cases, (I) when S_t is an axisymmetric torus, and, (II) when S_t is an axisymmetric sphere (there are no other possibilities). Take into account that all the calculations below are made on $(\Omega; g(t), K(t))$ in particular that $A(S_t)$ denotes the $g(t)$-area of S_t, i.e. $A(S_t) = A_{g(t)}(S_t)$.

(I) S_t *is an axisymmetric torus.* Choosing $\alpha = 1$ in the stability inequality (15) we obtain

$$\int_{S_t} |K|^2 dA \leq \int_{S_t} k^2 dA$$

and from (2) we get

$$(8\pi)^2|J(S_t)|^2 = |\int_{S_t} K(\xi,\varsigma)dA|^2 \leq \frac{\ell_0^2 A_1}{(2\pi)^2}\int_{S_t}|K|^2 dA$$

From these two inequalities and the bounds mentioned above we obtain

$$(8\pi)^2|J(S_t)|^2 \leq \frac{A_1^2\ell_0^2k_0^2}{(2\pi)^2}\ t^4$$

Now, if t_0, $(t_0 < \bar{t}_0)$ is small enough we obviously have

$$A_0 \geq \frac{A_1\ell_0 k_0\ t_0^2}{2\pi}$$

Putting all the inequalities together we obtain for any $t \leq t_0$ and $\Lambda_0 > 0$

$$A(S_t) \geq A_0 \geq \frac{A_0\ell_0k_0\ t_0^2}{2\pi} \geq 8\pi|J(S_t)| \geq 8\pi|J(S_t)| - \Lambda_0 t^4$$

(II) *S_t is an axisymmetric sphere.* Again, take into account that all the calculations below are made on $(\Omega; g(t), K(t))$. Let $\{\theta,\varphi\}$ be the areal-coordinates of the sphere S_t and let σ be as in (16). We would like to obtain first uniform upper and lower bounds for the function $\sigma(\theta)$ $(\theta \in [0,2\pi])$ where by uniform we mean independently on θ and independent also on the stable surface S_t $(t_0 \leq \bar{t}_0$, $\bar{t}_0$ as before), although dependent on ρ_0, μ_0 and μ_1. Observe that because any axisymmetric sphere has two poles then S_t intersects $\{\mathcal{P}\}$ at two points. By Corollary 3 if S_t intersects $\mathcal{T}_{g(t)}(\mathcal{P},\epsilon_2)$ it does so in discs D on which, in polar coordinates, we have the bounds

$$|\ell(s) - 2\pi s| \leq c_0 s^2, \tag{23}$$

$$|A(s) - \pi s^2| \leq c_1 s^3 \tag{24}$$

where we are using the notation for $\ell(s), A(s)$ as explained in Section 2.3. The equations are valid on $0 < s < s_D$ with $s_D \in [s_0, s_1]$ and the constants s_0, s_1, c_0, c_1 depend only on ρ_0, μ_0 and μ_1. Moreover we have (suppose without loss of generality that $\theta = 0$ is the pole of the disc),

$$\ell(s(\theta)) = 2\pi e^{\frac{\sigma(\theta)}{2}}\sin\theta, \tag{25}$$

$$A(s(\theta)) = \frac{A(S_t)}{2}(1-\cos\theta) \tag{26}$$

Combining (25) and (26) we get

$$e^{\sigma(\theta(s))} = \frac{A(s)}{\ell^2(s)}(A(S_t) - A(s))$$

To estimate the right hand side of this expression we will use

$$\frac{1}{4\pi}\left(\frac{1-c_1 s/\pi}{1+c_0 s/\pi}\right)\leq \frac{A(s)}{\ell^2(s)}\leq \frac{1}{4\pi}\left(\frac{1+c_1 s/\pi}{1-c_0 s/\pi}\right)$$

obtained from the inequalities (23)-(24), and,

$$A_0\leq A(S_t)\leq A_1$$

From them one easily shows that there is $s_2(\rho_0,\mu_0,\mu_1)$, with $s_2\leq s_0$, such that for any $s\in(0,s_2]$ we have (the coarse) bounds

$$\frac{A_0}{16\pi}\leq e^{\sigma}\leq \frac{A_1}{2\pi} \tag{27}$$

It follows that there is some uniform $\epsilon_3(\rho_0,\mu_0,\mu_2)$ with $\epsilon_3\leq\epsilon_2$ such that for any S_t, $|\sigma|$ is uniformly bounded on $S_t\cap\mathcal{T}_{g(t)}(\mathcal{P},\epsilon_3)$.

Moreover, combining (23)-(26) we obtain that there is θ_0 such for any point $q\in S_t$ outside $\mathcal{T}_{g(t)}(\mathcal{P},\epsilon_3)$ we have either $|\theta(q)|\geq\theta_0$ or $|\pi-\theta(q)|\geq\theta_0$. On the other hand there are uniform upper and lower bounds for the length $\ell(q)$ of the axisymmetric circles (orbits) passing through any point $q\in S_t$ outside $\mathcal{T}_{g(t)}(\mathcal{P},\epsilon_3)$. Because of these two facts and the expression $e^{\frac{\sigma(\theta(q))}{2}}=\frac{\ell(q)}{2\pi\sin\theta(q)}$ we deduce that there is a uniform bound for $|\sigma(\theta(q))|$ at any point $q\in S_t$ outside the tubular neighborhood $\mathcal{T}_{g(t)}(\mathcal{P},\epsilon_3)$. We thus obtain a uniform bound for $|\sigma|$ on any axisymmetric stable minimal sphere S_t with $A(S_t)\leq A_1$ as desired.

Now, following [6] (see also [7]) one can use the stability inequality to deduce[5] for axisymmetric spheres S_t the following crucial inequality

$$A(S)\geq 4\pi e^{\frac{\overline{\mathcal{M}}-8}{8}} \tag{28}$$

where $\overline{\mathcal{M}}$ is defined by

$$\overline{\mathcal{M}}=\frac{1}{2\pi}\int_S(\sigma'^2+4\sigma+\frac{\omega'^2}{\eta^2})\sin\theta d\theta d\phi-\frac{1}{2}e^{2c}\int_S k^2e^{-\sigma}\sin\theta d\theta d\phi$$

In this expression $e^c=A(S)/4\pi$, $\eta=e^{\sigma}\sin^2\theta$, the prime in σ' and ω' are their θ-derivative and $\omega=\omega(\theta)$ is a smooth function defined through

$$\frac{d\,\omega}{d\,\theta}=\frac{A(S)}{2\pi}K(\xi,\varsigma)\sin\theta$$

---

[5] Use (15)-(29)-(31)-(32) in [6] and instead of (30) in [6] use the constraint equation $R=|K|-k^2$. Finally note that the equation (2) for the angular momentum is still valid even if the data set is not maximal. Indeed the Komar angular momentum can be intrinsically defined as (5) in [7] which coincides with (2).

A direct computation shows $J(S) = (\omega(\pi) - \omega(0))/8$. On the other hand it is proved in [6], [1] that, if we let

$$\mathcal{M} = \frac{1}{2\pi}\int_S (\sigma'^2 + 4\sigma + \frac{\omega'^2}{\eta^2})\sin\theta d\theta d\phi$$

then

$$4\pi e^{\frac{\mathcal{M}-8}{8}} \geq 8\pi|J(S)| \tag{29}$$

Using this in (28) we obtain

$$e^{\left[\frac{e^{2c}}{16}\int_S k^2 e^{-\sigma}\sin\theta d\theta d\phi\right]}A(S) \geq 8\pi|J(S)| \tag{30}$$

We will use this now for $S = S_t$ on $(\Omega; g(t), K(t))$. Using the bounds $A(S_t) \leq A_1$, $|k| = |k(t)| \leq k_0 t^2$ and (27) one obtains

$$\left(\frac{A(S_t)}{16\pi}\right)^2 \int_{S_t} k^2 e^{-\sigma}\sin\theta d\theta d\phi \leq \frac{A_1^2 k_0^2}{16 A_0}t^4$$

Let $x = \frac{A_1^2 k_0^2}{16A_0}t^4$. Then if t_0 is suficiently small we have $0 < x \leq \frac{A_1^2 k_0^2}{16A_0}t_0^4 \leq 1$ and therefore $e^x \leq 1 + 2x$. Using this into (30) and using once more the bound $A(S_t) \leq A_1$ we obtain

$$A(S_t) \geq 8\pi|J(S_t)| - \frac{A_1^3 k_0^2}{16A_0}t^4$$

The claim follows by defining $\Lambda_0 = \frac{A_1^3 k_0^2}{16A_0}$. □

We are ready for the proof of Theorem 1. We recall first the setup of the proof. We assume by contradiction that there is an asymptotically flat data set $(\Sigma; g, K)$ on which there is an extreme Kerr-throat sphere. Then, as was explained, one can always consider a data set $(\Omega; g, K)$ (constructed from the data $(\Sigma; g, K)$) where $(\Omega; g)$ is a compact manifold with strictly mean convex boundary and having an extreme Kerr-throat sphere S_H in its interior. Moreover there exists a positive solution N_0 to the Lapse equation on Ω which is not proportional to α_T over the extreme sphere. The data $(\Omega; g, K)$ is embedded in a space-time and we consider its evolution under the vector field V as in (22) which gives us an axisymmetric flow $(\Omega; g(t), K(t))$.

Proof of Theorem 1. Let t_0 and Λ_0 be as in Proposition 5 (with $A_1 = A_{g(0)}(S_H)$). Let t be a time in $(0, t_0)$. Below we will work on $(\Omega; g(t), K(t))$. Therefore keep in mind that all the quantities, in particular areas, are found from $(g(t), K(t))$.

First we observe that the infimum of the areas of all the surfaces isotopic to S_H is non-zero. Indeed for any surface S' isotopic to S_H we have

$$0 < |J(S_H)| = |J(S')| = \frac{1}{8\pi}|\int_{S'} K(\xi, \varsigma)dA| \leq \frac{1}{8\pi}\|K\|_{L^\infty}\|\xi\|_{L^\infty}A(S')$$

Then, following [8] (Theorems 1 and 1') there is a sequence of surfaces $\{S'_l\}$, with each S'_l isotopic to S_H, converging in measure to $n_1S_1+\ldots+n_kS_k$, where $\{S_1,\ldots,S_k\}$ is a set of compact and embedded surfaces[6][7]. Moreover the infimum of the areas of all the surfaces isotopic to S_H is equal to $n_1A(S_1)+\ldots+n_kA(S_k)$. In particular $A(S_i)\leq A(S_H)\leq A_1$ (this upper bound is needed to apply later Proposition 5). If one of the surfaces, say S_i, is orientable, then it is stable and therefore axisymmetric (see Section 2.5). In such case there are $n_i^+\geq 0$, $n_i^-\geq 0$ with $n_i=n_i^++n_i^-$, indicating how many times the sequence $\{S'_l\}$ "wraps around" the oriented S_i with one orientation and how many with the opposite orientation. Precisely, for any two-form χ supported on a small neighborhood of the oriented surface S_i, we have

$$\lim\int_{S'_l}\chi=(n_i^+-n_i^-)\int_{S_i}\chi$$

If on the other hand one of the surfaces, say S_i, is non-orientable then for any two-form supported in a small neighborhood S_i we have

$$\lim\int_{S'_l}\chi=0 \tag{31}$$

Note that if all the S_i's were non-orientable, then using (31) with $\chi=*K(\xi,-)$, we would get $J(S_H)=J(S'_l)=\lim J(S'_l)=0$ which is not possible. We deduce that at least one of S_i's has to be orientable (this fact is not essential). Let us order the surfaces in such a way that $\{S_1,\ldots,S_j\}$, $j\geq 1$ are the orientable (and oriented) and $\{S_{j+1},\ldots,S_k\}$ are the non-orientable. We have

$$\begin{aligned}|J(S_H)|=\lim|J(S'_l)|=|\sum_{i=1}^{i=j}(n_i^+-n_i^-)J(S_i)|&\leq\sum_{i=1}^{i=j}n_i|J(S_i)|\leq\sum_{i=1}^{i=j}\frac{n_i}{8\pi}A(S_i)+O(t^4)\\&\leq\frac{1}{8\pi}A(S_H)+O(t^4)\leq|J(S_H)|-\frac{A''_{N_0}(S_H)}{16\pi}t^2+O(t^3).\end{aligned}$$

where $A''_{N_0}(S_H)>0$ and where to obtain the inequality between the fourth and fifth terms we have used the Proposition 5 and to obtain the inequality between the sixth and seventh terms we have used Proposition 3. We obtained thus a contradiction for short times. This finishes the proof of Theorem 1. □

3.3 Proof of Theorem 2.

In the following sections the reader may benefit from the following "quotient" viewpoint on the geometry of (Σ,g), where (Σ,g), as in the hypothesis of Theorem 2, is axisymmetric and diffeomorphic to $S^2\times\mathbb{R}$.

Recall that the group $U(1)$ acts by isometries and that the set of fixed points consist of two connected and complete one-dimensional manifolds (the axes), and therefore each diffeomoprhic to $\mathbb{R}$. The quotient of Σ by the action, denoted by $\tilde{\Sigma}$ is diffeomorphic to $[0,1]\times\mathbb{R}$, where $\{\mathcal{S}\} := \{0\}\times\mathbb{R}$ and $\{\mathcal{N}\} := \{1\}\times\mathbb{R}$ are the pair of "South" and "North" axis. The set of both axis will be denoted as before by $\{\mathcal{P}\} = \{\mathcal{N}\}\cup\{\mathcal{S}\}$ and the topological interior by $\tilde{\Sigma}^\circ := \tilde{\Sigma}\setminus\{\mathcal{P}\} = \tilde{\Sigma}\setminus\partial\tilde{\Sigma}$. We denote the projection by Π (in particular $\Pi(\Sigma) = \tilde{\Sigma}$). Denote by λ^2 the square norm of the axisymmetric Killing field and let $\tilde{g}$ be the two dimensional quotient metric on $\tilde{\Sigma}^\circ$, namely, if $\tilde{w} = \Pi(w)$ and $\tilde{v} = \Pi(v)$ with w, v tangent vectors at $p\in\Sigma\setminus\{\mathcal{P}\}$, then at $\tilde{p} = \Pi(p)$ we have

---

[6] A very accurate description of the relation between the sequence $\{S'_l\}$ and the surfaces $S_1,\ldots,S_k$ is given in Remark (3.27) of [8].

[7] Namely for every function f on Ω we have $\lim\int_{\bar{S}_l}fdA=\sum n_i\int_{S_i}fdA$.

$$\tilde{g}(\tilde{w},\tilde{v}) = g(w,v) - \frac{g(\xi,w)g(\xi,v)}{\lambda^2}$$

The metric $\tilde{g}$ extends smoothly to $\tilde{\Sigma}$. Every axisymmetric sphere S projects into a one-dimensional manifold diffeomorphic to $[0,1]$ starting and ending $\tilde{g}$-perpendicularly to the axis (see Figure 4). An axisymmetric sphere is contractible iff the projection starts and ends at the same axis. Axisymmetric torus project into closed curves inside $\tilde{\Sigma}^\circ$ and are therefore contractible in Σ. Besides spheres and tori, there are no more orientable axisymmetric boundary-less surfaces. Axisymmetric discs project into one-dimensional manifolds diffemorphic to $[0,1]$, starting $\tilde{g}$-perpendicularly at an axis and ending at an interior point. We will use the notation $\beta^{\mathcal{N},\mathcal{S}}$ for projected axisymmetric spheres starting at $\{\mathcal{N}\}$ and ending at $\{\mathcal{S}\}$, and $\beta^{\mathcal{N}}(\tilde{p})$ (resp. $\beta^{\mathcal{S}}(\tilde{p})$) for projected discs starting at $\tilde{p}$ and ending at $\{\mathcal{N}\}$ (resp. $\{\mathcal{S}\}$). All this is shown in Figure 4.

Let $\beta(\tilde{s})$ be a curve in $\tilde{\Sigma}$ parametrized with respect to $\tilde{g}$ arc-length, then

$$A(\Pi^{-1}(\beta)) = 2\pi\int_{s_0}^{s_1}\lambda(\beta(\tilde{s}))\,d\tilde{s}$$

In other words the area is equal to the length of β with respect to the conformal metric $\bar{\tilde{g}} = (2\pi\lambda)^2\tilde{g}$. In particular axisymmetric minimal surfaces (which minimize area locally) in Σ correspond to $\bar{\tilde{g}}$-geodesics in $\tilde{\Sigma}^\circ$. Thus, the well known property that if two minimal surfaces are tangent at a point then they coincide is direct in the context of axisymmetric minimal surfaces. This fact will be useful later.

The hypothesis of Theorem 2 translates into the following two conditions on the quotient manifold:

C1. There is a $\bar{\tilde{g}}$-geodesic $\gamma_H^{\mathcal{N},\mathcal{S}}$ of $\bar{\tilde{g}}$-length $A_H := 8\pi|J|$.

C2. For any curve $\beta^{\mathcal{N},\mathcal{S}}$, its $\bar{\tilde{g}}$-length is greater or equal than A_H.

(we will use the notation A_H for $8\pi|J|$ from now on). The curve $\gamma_H^{\mathcal{N},\mathcal{S}}$ divides $\tilde{\Sigma}$ into two smooth manifolds $\tilde{\Sigma}_1$ and $\tilde{\Sigma}_2$ each diffeomorphic to $[0,1]\times\mathbb{R}^+$ ($\mathbb{R}^+ = [0,\infty)$). Denote $\Sigma_i = \Pi^{-1}(\tilde{\Sigma}_i)$. At a point $\tilde{p}\in\tilde{\Sigma}_i$, with i either 1 or 2, define

$$\underline{A}_i(\tilde{p}) = \inf\{\text{length}_{\bar{\tilde{g}}}(\beta^{\mathcal{N},\mathcal{S}}(\tilde{p})),\ \beta^{\mathcal{N},\mathcal{S}}(\tilde{p})\subset\tilde{\Sigma}_i\},$$
$$\underline{A}_i^{\mathcal{N}}(\tilde{p}) = \inf\{\text{length}_{\bar{\tilde{g}}}(\beta^{\mathcal{N}}(\tilde{p})),\ \beta^{\mathcal{N}}(\tilde{p})\subset\tilde{\Sigma}_i\}$$



Figure 4

and similarly for $\underline{A}_i^{\mathcal{S}}(\tilde p)$. We also define

$$\underline{A}(\tilde p) = \{\text{length}_{\tilde{\bar g}}(\beta^{\mathcal{N},\mathcal{S}}(\tilde p)),\ \beta^{\mathcal{N},\mathcal{S}}(\tilde p)\subset\tilde\Sigma\} \tag{32}$$

In the original space, $\underline{A}_i(\tilde p)$ is just the infimum of the areas of axisymmetric spheres in Σ_i containing the orbit $\Pi^{-1}(\tilde p)$. Similarly $\underline{A}_i^{\mathcal{N}}(\tilde p)$ (resp. $\underline{A}_i^{\mathcal{N}}(\tilde p)$) is the infimum of the areas of axisymmetric discs in Σ_i intersecting $\{\mathcal{N}\}$ (resp. $\{\mathcal{S}\}$) and with boundary $\Pi^{-1}(\tilde p)$. These quantities are irrelevant outside axisymmetry. By **C2** we have $\underline{A}_i(\tilde p)\geq A_H$ for all $\tilde p$.

The use of the quotient picture has considerable advantages but also shortcomings. As a general rule the analysis away from the axis is more conveniently done in the quotient geometry. At the axis however the metric $\tilde{\bar g}$ is singular and it is better to stick to the original space. For this reasons we will keep a mixed usage.

We state now a basic Proposition concerning area-minimizing sequences of discs that will be required to prove Lemma 1 on which the proof of Theorem 2 relies. To avoid further delays for the proof of Theorem 2 we postpone the proof of the Proposition until the Appendix. We use the notation $D^{\mathcal{N}}(C)$ and $D^{\mathcal{S}}(C)$ as in Section 2.3 for axisymmetric discs intersecting $\{\mathcal{N}\}$ and $\{\mathcal{S}\}$ respectively.

Proposition 6 *Let i be 1 or 2. Let $\Pi(C)=\tilde p\in\tilde\Sigma_i^\circ$. Then $\underline{A}_i^{\mathcal{N}}(\tilde p)$ is equal to either*

D1. *the area of a stable minimal disc $D^{\mathcal{N}}(C)=\Pi^{-1}(\gamma^{\mathcal{N}}(\tilde p))$, or,*

D2. *the area of a stable minimal disc $D^{\mathcal{S}}(C)=\Pi^{-1}(\gamma^{\mathcal{S}}(\tilde p))$ plus A_H.*

Moreover there is an area minimizing sequence of discs $\{D_j^{\mathcal{N}}(C)\}=\{\Pi^{-1}(\beta_j^{\mathcal{N}}(\tilde p))\}$ converging (in measure) to either

$D^{\mathcal{N}}(C)=\Pi^{-1}(\gamma^{\mathcal{N}}(\tilde p))$ in case **D1** *holds, or,*

$D^{\mathcal{S}}(C)\cup S_H=\Pi^{-1}(\gamma^{\mathcal{S}}(\tilde p))\cup\Pi^{-1}(\gamma_H^{\mathcal{N},\mathcal{S}})$ in case **D2** *holds.*

A similar statement holds for $\underline{A}_i^{\mathcal{S}}(\tilde p)$ by changing $\mathcal{N}\to\mathcal{S}$ and $\mathcal{S}\to\mathcal{N}$.

Lemma 1 *Let i be 1 or 2. If for all $\tilde{p} \in \tilde{\Sigma}_i$ it is $\underline{A}_i(\tilde{p}) = A_H$ then $(\Sigma_i; g, K)$ is half of the extreme Kerr throat of angular momentum $J| = A_H/8\pi$.*

Proof: Before we start recall that any stable minimal surface S with $A(S) = 8\pi|J(S)|$ is an extreme Kerr-throat sphere [6]. We prove first that Σ_i is foliated by extreme Kerr-throat spheres, or, in the quotient space, that $\tilde{\Sigma}_i$ is foliated by $\bar{\tilde{g}}$-geodesics of length A_H starting $\tilde{g}$-perpendicularly to $\{\mathcal{N}\}$ and ending $\tilde{g}$-perpendicularly to $\{\mathcal{S}\}$. In this first part of the proof we work in the quotient space. Let $\tilde{p} \in \tilde{\Sigma}_i^\circ$. By hypothesis $\underline{A}_i(\tilde{p}) = A_H$. Therefore, by Proposition 6, $\underline{A}_i^{\mathcal{N}}(\tilde{p})$ and $\underline{A}_i^{\mathcal{S}}(\tilde{p})$ are realized $\bar{\tilde{g}}$-geodesics $\gamma^{\mathcal{N}}(\tilde{p})$ and $\gamma^{\mathcal{S}}(\tilde{p})$ respectively (case **D1** holds) and moreover the sum of their $\bar{\tilde{g}}$-lengths is A_H. If $\gamma^{\mathcal{N}}(\tilde{p})$ and $\gamma^{\mathcal{S}}(\tilde{p})$ do not have the same tangent line at $\tilde{p}$ then their union can be rounded up at the vertex $\tilde{p}$ to a curve with $\bar{\tilde{g}}$-length less than A_H which violates **C2**. Thus $\gamma^{\mathcal{N},\mathcal{S}}(\tilde{p}) := \gamma^{\mathcal{N}}(\tilde{p}) \cup \gamma^{\mathcal{S}}(\tilde{p})$ is a $\bar{\tilde{g}}$-geodesic of length A_H or, the same, $\Pi^{-1}(\gamma^{\mathcal{N}}(\tilde{p}) \cup \gamma^{\mathcal{S}}(\tilde{p}))$ is an extreme Kerr-throat sphere. We claim that for any $\tilde{p}_1 \neq \tilde{p}_2$ in $\tilde{\Sigma}_i^\circ$ the geodesics $\gamma^{\mathcal{N},\mathcal{S}}(\tilde{p}_1)$ and $\gamma^{\mathcal{N},\mathcal{S}}(\tilde{p}_2)$ must be either equal or disjoint. Indeed if they are not disjoint then they cannot be tangent other the geodesics must coincide, and if they intersect transversely then one can again construct easily a curve $\beta^{\mathcal{N},\mathcal{S}}(\tilde{p})$ of $\bar{\tilde{g}}$-length less than A_H which violates again **C2**. The set $\{\gamma^{\mathcal{N},\mathcal{S}}(\tilde{p}), \tilde{p} \in \Sigma_i^\circ\}$ is thus the desired foliation of $\tilde{\Sigma}_i$ by $\bar{\tilde{g}}$-geodesics of length A_H and $\{\Pi^{-1}(\gamma^{\mathcal{N},\mathcal{S}}(\tilde{p})), \tilde{p} \in \Sigma_i^\circ\}$ is the desired foliation of Σ_i by extreme Kerr-throat spheres.

So far we have a foliation by extreme Kerr-throat spheres. From this information we want to deduce that there are coordinates $\{(r, \theta, \varphi)\}$ on Σ_i on which the metric g has exactly the expression (4). Let r be a smooth function, constant along the leaves of the foliation and with non-zero gradient everywhere. The flow induced by the vector field

$$Y := \frac{\nabla^i r}{|\nabla r|^2}$$

takes leaves into leaves ($dr(Y) = 1$) (and indeed orbits into orbits because, can be seen, Y is $U(1)$-invariant). Let $\{(\theta, \varphi)\}$ be areal coordinates on S_H. Extend them to all Σ_i by Lie dragging, namely define them by imposing $Y(\theta) = 0,\ Y(\varphi) = 0$. In this way $\{(r, \theta, \varphi)\}$ are coordinates on Σ_i. Let $h_{AB}(\bar{r})$ be the metric components of the two-metric induced on the leaf $\{r = \bar{r}\}$ in the coordinates $\{(\theta, \varphi)\}$. Then we have

$$\partial_r h_{AB} = 0 \tag{33}$$

because every leaf is totally geodesic. Thus $h_{AB}(r) = h_{AB}(0)$ which is the metric of S_H in areal coordinates, namely (from (4))

$$h(0) = \frac{4|J| \sin^2\theta}{1 + \cos^2\theta} d\varphi^2 + |J|(1 + \cos^2\theta) d\theta^2$$

By (33) the first and second variation of the area of the leaves along Y is zero. We deduce that at every leaf we must have

$$\partial_r = Y = c(r) \alpha_T \varsigma$$

where ς is a g-unit normal field to the leaf. It follows that one can redefine r to have $\partial_r = \alpha_T \varsigma$ over every leaf. As $< \partial_r, \partial_\theta >_g = < \partial_r, \partial_\varphi >_g = 0$, the metric g takes in these coordinates the form

$$g = \alpha_T^2 dr^2 + h_{AB}(0)$$

which is (4). That the second fundamental form takes the form (5) is a consequence of the fact that, by the uniqueness mentioned before, K takes the form of (5) over any extreme Kerr-throat sphere. □

We are ready to prove Theorem 2.

Proof of Theorem 2. As before S_H divides Σ into Σ_1 and Σ_2 (both including S_H). Let $\bar{A}_i = \sup\{\underline{A}_i(\tilde{p}), \tilde{p} \in \Sigma_i\}$, for $i = 1, 2$. If $\bar{A}_i = A_H$ for $i = 1, 2$ then by Lemma 1 the sate has to be the extreme Kerr throat. On the other hand suppose that one (but not both) of the $\bar{A}_i$'s is equal to A_H then one can easily make a doubling to construct a state with an extreme Kerr-sphere S'_H, and two sectors Σ'_1, Σ'_2 with $\bar{A}_i > A_H$ for $i = 1, 2$. This shows that to prove Theorem 2 it is enough to assume that $\bar{A}_i > A_H$ for $i = 1, 2$ and obtain a contradiction. We will assume that from now on.

Let $\tilde{p}_1 \in \tilde{\Sigma}_1^\circ$ and $\tilde{p}_2 \in \tilde{\Sigma}_2^\circ$ be points such that $\underline{A}_i(\tilde{p}_i) > A_H$ for $i = 1, 2$. Now observe that the set of points $\mathcal{E} = \{\tilde{p} \in \tilde{\Sigma}^\circ, \underline{A}(\tilde{p}) = A_H\}$ ($\underline{A}(\tilde{p})$ is defined in (32) is

(i) a closed set, and,

(ii) a union of projected extreme Kerr-throat spheres.

That $\mathcal{E}$ is closed follows from the fact that $\underline{A}(\tilde{p})$ is continuous with respect to $\tilde{p}$. That $\mathcal{E}$ is a union of projected extreme Kerr-throat spheres follows from the fact, shown inside the proof of Lemma 1, that if $\underline{A}(\tilde{p}) = A_H$ then $\tilde{p}$ lies in a projected extreme Kerr-throat sphere.

Let S_i, $i = 1, 2$ be spheres (with poles in $\{\mathcal{N}\}$ and $\{\mathcal{S}\}$ denoted by $\mathcal{N}_i$ and $\mathcal{S}_i$) such that

1. $S_i \subset (\Sigma_i \setminus \Pi^{-1}(\mathcal{E})$, and,
2. Let $R > 0$ such that the space $(B_{\frac{g}{R^2}}(\mathcal{N}_i, 2), \frac{g}{R^2})$ is ϵ_0-close[8] in C^2 to the flat metric in $B(o, 2) \subset \mathbb{R}^3$ where as in Section 2.5, $B(o, 2)$ is the Euclidean ball of radius 2 and ϵ_0 is as in Proposition 2. Then $S_i \cap B_{\frac{g}{R^2}}(\mathcal{N}_i, L)$ is a disc as in *item 3* of Corollary 3.

Let Ω be the region enclosed by S_1 and S_2 (including them). Let $\delta > 0$ be such that

1'. $\mathcal{T}(S_i, 4\delta) \subset (\Sigma \setminus \Pi^{-1}(\mathcal{E}))$, and,

2'. There are g-Gaussian coordinates on $\mathcal{T}(S_i, 4\delta)$.

Following Proposition 4, let N_0 be a positive solution of the Lapse equation in

$$\Omega_{4\delta} := \Omega \cup \mathcal{T}(S_1, 4\delta) \cup \mathcal{T}(S_2, 4\delta)$$

---

[8] To be precise in the coordinates $\{x^1, x^2, x^3\}$ as in Section 2.4.

which is not proportional to α_T on S_H. Evolve

$$\Omega_{3\delta} := \Omega \cup \mathcal{T}(S_1, 3\delta) \cup \mathcal{T}(S_2, 3\delta)$$

along the space following the vector field V as in (22) thus obtaining a flow

$$(\Omega_{3\delta}; g(t), K(t); N(t), X(t))$$

on the fixed manifold $\Omega_{3\delta}$, for $t \in [0, t_0]$.

Consider now a continuous path $S_i(t)$, $i = 1, 2$ of spheres (with poles on $\{\mathcal{N}\}$ and $\{\mathcal{S}\}$ denoted by $\mathcal{N}_i(t)$ and $\mathcal{S}_i(t)$) such that

1". $S_i(0) = S_i$, $i = 1, 2$, and,

2". The space $(B_{\frac{g(t)}{R_1^2}}(\mathcal{N}_i(t), 2), \frac{g(t)}{R_1^2})$ is ϵ_0-close to in C^2 to the flat metric in $B(o, 2)$, where $R_1 = \min\{R, \delta/2\}$, and $S_i(t) \cap B_{\frac{g(t)}{R_1^2}}(\mathcal{N}_i(t), L)$ is a disc as in *item 3* of Corollary 3. Moreover the same we assume occurs at the south poles $\mathcal{S}_i(t)$.

Let t_0 be smaller if necessary such that for any $t \in [0, t_0]$ there are $g(t)$-Gaussian coordinates in $\mathcal{T}_{g(t)}(S_i(t), 2\delta)$. Define $\Omega(t)$ the region enclosed by $S_1(t)$ and $S_2(t)$ and

$$\Omega_{2\delta}(t) = \Omega(t) \cup \mathcal{T}_{g(t)}(S_i(t), 2\delta) \cup \mathcal{T}_{g(t)}(S_2(t), 2\delta)$$

Now, instead of $g(t)$ we will consider a modified flow of axisymmetric metrics $g^*(t)$ on $\Omega_{2\delta}(t)$ with the following properties for every t

1. $g^*(t) = g(t)$ on $\Omega(t)$,
2. $g^*(t) \geq g(t)$ on $\Omega_{2\delta}(t)$,
3. the g^*-mean curvature of the boundary of the region $\Omega_{2\delta}(t)$ is strictly positive in the outward direction, namely the boundary is strictly mean convex.

The flow of metrics $g^*(t)$ can be explicitly given for instance as follows. On everyone of the two connected components of $\Omega_{2\delta}(t) \setminus \Omega(t)$ we write the metric $g(t)$ in $g(t)$-Gaussian coordinates as $g(t) = dr^2 + h_i(t)$, $i = 1, 2$. Then make

$$g^*(t) := \begin{cases} g(t) & \text{on} \quad \Omega(t), \\ dr^2 + f^2(r) h_i(t), & \text{on} \quad \Omega_{2\delta}(t) \setminus \Omega(t), \ (r > 0), \end{cases}$$

where $f(r)$ is the scalar and time independent function

$$f(r) = 1 + e^{-\frac{1}{r} + \frac{1}{2\delta + \epsilon - r}}$$

for $\epsilon > 0$ small enough making the boundary strictly convex.

Now, at time zero we have $\ddot{A}(S_H) < 0$ and therefore for any $t < t_0$ (t_0 smaller if necessary) we have $A_{g(t)}(S_H) < A_H$. By [8] (Theorems 1 and 1') there is, for every $0 < t \leq t_0$, a $g^*(t)$-stable, axisymmetric and area minimizing sphere S_t inside $\Omega_{2\delta}(t)$ of area less than A_H. We claim that (making t_0 smaller if necessary) S_t must lie inside $\Omega(t)$ where the metric $g^*(t)$ is equal to $g(t)$ and therefore showing the "barrier"-like character of the $S_i(t)$'s and the $g(t)$- stability of S_t. We show this now. Let $\Gamma > 0$ be such that for every sphere S (with poles in $\{\mathcal{N}\}$ and $\{\mathcal{S}\}$) intersecting $\Omega_{2\delta} \setminus (\Omega \cup \mathcal{T}(\{\mathcal{P}\}, R_1 L))$ has g-area greater or equal than $A_H + \Gamma$. By continuity, and making t_0 smaller if necessary, we have that for every $t \in [0, t_0]$ and every sphere S (with poles in $\{\mathcal{N}\}$ and $\{\mathcal{S}\}$) intersecting $\Omega_{2\delta}(t) \setminus (\Omega(t) \cup \mathcal{T}_{g(t)}(\{\mathcal{P}\}, R_1 L))$ has $g^*(t)$-area greater or equal than $A_H + \Gamma/2$. Now, let S_t be the area minimizing sphere of $g^*(t)$-area less than A_H as considered before. If S_t lies not in $\Omega(t)$ then, being an sphere with poles in $\{\mathcal{N}\}$ and $\{\mathcal{S}\}$ and of $g^*(t)$-area less than A_H it must necessarily intersect the discs $S_i(t) \cap B_{g(t)/R_1^2}(\mathcal{N}_i(t), L)$ and $S_i(t) \cap B_{g(t)/R_1^2}(\mathcal{S}_i(t), L)$ which violates Proposition 2. Thus $S_t \subset \Omega(t)$.

Finally in the same way as in Proposition 5, one can consider t_0 smaller if necessary and $\Lambda_0 > 0$ such that for any $t \in (0, t_0]$ we have $A(S_t) = A_{g(t)}(S_t) \geq 8\pi|J| - \Lambda_0 t^4$. On the other hand $A(S_t) = A_{g(t)}(S_t) \leq A_H + \ddot{A}(S_H)t^2/2 + O(t^3)$. This is a contradiction as $A_H = 8\pi|J|$ and $\ddot{A}(S_H) < 0$. $\square$

4 Appendix.

Proof of Proposition 1. We prove first *item 1* which is true for any value of L chosen between $(0,1)$ (indeed we just reproduce here the classical proof). Let D be a minimal disc with boundary C. Say $z|_C = z_C$. Then the function $z - z_C$ is harmonic on D and $(z - z_C)|_C = 0$. It follows that $z = z_C$ all over D and therefore that D is the disc enclosed by C in the plane $\{z = z_C\}$. Another proof, best suited for extensions, can be obtained along the following lines (we just provide the sketch). Let $D(C)$ be the disc enclosed by C in the plane $\{z = z_C\}$. Let $o(C) = (0, 0, z_C)$ be its center and $R(C)$ its radius. Let $B^+ = B(o(C), R(C)) \cap \{z > z_C\}$ and $B^- = B(o(C), R(C)) \cap \{z < z_C\}$. Now, the disc $D(C)$ is minimal and, by a direct inspection of the stability operator, also strictly stable. Using the strict stability one can construct a smooth foliation of B^+ by discs $\{D'(C)\}$, each with boundary C, and strictly convex (in the direction of increasing z) and similarly for B^-. In addition one can construct a foliation of $B(o,1) \setminus B(o(C), R(C))$ by round spheres $\{S'\}$. We have thus a foliation of $B(o,1) \setminus D(C)$ by strictly convex surfaces, acting as barriers, and preventing the existence of any other minimal disc with boundary C inside $B(0,1)$.

We prove now *item 2*. First we prove that there are no stable surfaces in the class $\mathscr{S}_0$. This again is true for any value of L chosen in $(0,1)$. This can be proved as in *item 1* by showing that if there is one then the function z has to be constant on it. Onother proof, more independent of the flatness of the ambient space $\mathbb{R}^3$ and therefore best suited for extensions is the following. Assume again that there is one such surface. Then note that there is a cylinder $\{\rho = \rho_0\}$ where $\rho^2 = x^2 + y^2$, enclosing the surface and tangent to it at least in one orbit (a circle). Such cylinder has strictly mean convex boundary which implies that at the points of tangency the minimal surface must have positive mean curvature (in the outgoing direction from the axis) which is absurd. Note that instead of cylinders one could have used spheres to reach a similar conclusion.

We prove now that for some L appropriately chosen there are no stable surfaces in the class $\mathscr{S}_2$. This requires a bit more effort. Recall that the class $\mathscr{S}_2$ consist of axisymmetric cylinders with boundary in $B(o,1)$. From now on we let S be an stable axisymmetric cylinder with boundary in $\partial B(o,1)$. If $S\cap B(o,R)\neq\emptyset$ and $q\in S\cap B(o,R)$ we will denote by $[S\cap B(o,R)]^{c.c.}_q$ the connected component of $S\cap B(o,R)$ containing q. Following [5] Corollary 2.11 (pg. 79) we have that for any S such that $S\cap B(o,1/8)\neq\emptyset$ and $q\in S\cap B(o,1/8)$ then

$$\sup_{p\in[S\cap B(o,1/2)]^{c.c.}_q} |\Theta|^2(p)\leq 4c \tag{34}$$

where c is a universal constant (use in Corollary 2.11 $r_0=7/8$, $\sigma=1/2$ and that $B(o,1/4)\subset B(q,r_0-\sigma)$). Let $L_0=\min\{1/(4\sqrt{16c}),1/4\}$. Then from (34) we obtain that for any S such that $S\cap B(o,L_0/2)\neq\emptyset$ and $q\in S\cap B(o,L_0/2)$ we have

$$16L_0^2\sup_{p\in[S\cap B(o,L_0)]^{c.c.}_q} |\Theta|^2(p)\leq \frac{1}{16} \tag{35}$$

Let L be any number $0<L<L_0/8$. From (35) and Lemma 2.4 (pg. 74) in [5] we obtain that, for any surface S such that $S\cap B(o,L)\neq\emptyset$ and $q\in S\cap B(o,L)$ we have

1. $B_S(q,L_0/2)$ is a graph of a function u on a domain of $T_qS\subset\mathbb{R}^3$, where $B_S(q,L_0/2)$ is the intrinsic ball (in S with the induced metric) and radius $L_0/2$, and,
2. the connected component of $S\cap B(o,L_0/4)$ containing q, namely $[S\cap B(o,L_0/4)]^{c.c.}_q$, lies inside $B_S(1,L_0/2)$ (this follows easily from (2) in Lemma 2.4 in [5]).

For any surface S such that $S\cap B(o,L)\neq\emptyset$ let $\mathscr{C}=[S\cap B(o,L_0/4)]^{c.c}_q$ and thus $\mathscr{C}$ is a cylinder with boundary equal to two orbits, C_1 and C_2, in $\partial B(o,L_0/4)$. Denote by Π_{T_qS} the projection into $T_qS\subset\mathbb{R}^3$. Then $\Pi_{T_qS}(\mathscr{C})$ is an annulus with boundary components $\Pi_{T_qS}(C_1)$ and $\Pi_{T_q(S)}(C_2)$. Moreover $\Pi_{T_q(S)}C(q)$ ($C(q)$ is the orbit passing through q) encloses either $\Pi_{T_qS}(C_1)$ or $\Pi_{T_q(S)}(C_2)$. Say it encloses $\Pi_{T_q(S)}(C_2)$. Observe also that $\text{length}(\Pi_{T_qS}(C(q)))\leq\text{length}(C(q))\leq 2\pi L$ (as $q\in B(0,L)$). Let α be the curve in $\mathscr{C}$ with constant azimuthal angle φ and joining q to C_2 and observe that $\Pi_{T_qS}(\alpha)$ is a straight segment joining q to $\Pi_{T_qS}(C_2)$. Then because $|\nabla u|\leq 1$ we have $\text{length}(\alpha)\leq 2\text{length}(\Pi_{T_qS}(\alpha))\leq 2L$ (the last inequality is due to fact that the length of straight segment inside an ellipse with perimeter less than $2\pi L$ has length less than L). But on the other hand $q\in B(o,L)$ and $\mathscr{C}\subset\partial B(o,L_0/4)$ and therefore $\text{length}(\alpha)\geq L_0/4-L$. We conclude that there are no stable minimal surfaces S in the family $\mathscr{S}_2$ intersecting $B(o,L)$ if $L<L_0/20$. □

The proof of Proposition 2 is done in the same way as in Proposition 1 and will not be included here. Let us prove now Corollary 3.

Proof of Corollary 3: In several parts of the proof we will use the "quotient picture" as explained in Section 3.3.

Item 1. By homogeneous regularity there is $\epsilon_1(\rho_0, \mu_0, \mu_1)$ such that for any $0 < \epsilon \leq \epsilon_1$, the set

$$(\partial \mathcal{T}(\partial \Sigma, \epsilon)) \setminus \partial \Sigma,$$

is smooth and strictly convex with mean curvature greater or equal than $\mu_0/2$. Thus these surfaces act as barriers preventing the the existence of minimal surfaces S at a distance less than ϵ_1 from $\partial\Sigma$.

Item 2. By homogeneous regularity there is $\epsilon_4(\rho_0, \mu_0, \mu_1) > 0$ such that for any $0 < \epsilon < \epsilon_4$ the surface

$$(\partial \mathcal{T}(\{\mathcal{P}\}, \epsilon)) \cap (\Sigma \setminus \mathcal{T}(\partial \Sigma, \epsilon_1))$$

is smooth (with boundary) and of mean curvature greater or equal than one (in the outgoing direction from the axes $\{\mathcal{P}\}$). Note that as $\epsilon \to 0$ the mean curvature goes to infinity ($\sim 1/\epsilon$). In particular, because these surfaces act as barriers there are no minimal surfaces lying entirely inside

$$\mathcal{T}(\{\mathcal{P}\}, \epsilon_4) \cap (\Sigma \setminus \mathcal{T}(\partial \Sigma, \epsilon_1))$$

We will work now in the quotient and follow the notation of Section 3.3. By homogeneous regularity again, there is $R(\rho_0, \epsilon_1, \epsilon_4) > 0$ such that for any $\tilde{p}$ in $\tilde{\Sigma} \setminus \Pi\big(\mathcal{T}(\{\mathcal{P}\}, \epsilon_4) \cup \mathcal{T}(\partial\Sigma, \epsilon_1)\big)$ we have

$$B_{\bar{\bar{g}}}(\tilde{p}, R) \subset \left(\tilde{\Sigma} \setminus \Pi\big(\mathcal{T}(\{\mathcal{P}\}, \frac{\epsilon_4}{2}) \cup \mathcal{T}(\partial\Sigma, \frac{\epsilon_1}{2})\big)\right)$$

and moreover the metric $\frac{1}{R^2}\bar{\bar{g}}$ in

$$B_{\frac{1}{R^2}\bar{\bar{g}}}(\tilde{p}, 1) = B_{\bar{\bar{g}}}(\tilde{p}, R)$$

is sufficiently close in C^2 to the flat metric in $\mathbb{R}^2$ that every $\frac{1}{R^2}\bar{\bar{g}}$-geodesic (or, the same, every $\bar{\bar{g}}$-geodesic) passing through $\tilde{p}$ reaches the boundary of $B_{\frac{1}{R^2}\bar{\bar{g}}}(\tilde{p}, 1)$ and therefore has $\frac{1}{R^2}\bar{\bar{g}}$-length greater than 1 (or, the same, the $\bar{\bar{g}}$-length is greater or equal than R). Now, for every axisymmetric minimal surface S, $\Pi(S)$ is a $\bar{\bar{g}}$-geodesic and $A(S) = \text{length}_{\bar{\bar{g}}}(\Pi(S))$. Moreover by what was said before there is always a point $\tilde{p}$ of $\Pi(S)$ in $\tilde{\Sigma} \setminus \Pi\big(\mathcal{T}(\{\mathcal{P}\}, \epsilon_4) \cup \mathcal{T}(\partial\Sigma, \epsilon_1)\big)$. It follows that $A(S) \geq R$. The *item 2* follows by defining $A_0 := R$.

Item 3. By homogeneous regularity there exists $R_1(\rho_0, \epsilon_1, \epsilon_4) > 0$ such that for any $p \in \{\mathcal{P}\} \setminus \mathcal{T}(\partial\Sigma, \epsilon_1/2)$ the metric $\frac{1}{R_1^2} g$ in

$$B_{\frac{1}{R_1^2}}(p, 2)$$

is ϵ_0-close in C^2 to the flat metric in $\mathbb{R}^3$ where ϵ_0 is as in Proposition 2. Then we define

$$\epsilon_2 := L R_1$$

where L is as in Proposition 2. We show now that with this ϵ_2 we have all the properties that we desire for *item 3*. The cylinder (with boundary)

$$\big(\partial\mathcal{T}(\{\mathcal{P}\},\epsilon_2)\big)\cap B_g(p,R_1)=\big(\partial\mathcal{T}_{\frac{1}{R_1^2}g}(\{\mathcal{P}\},L)\big)\cap B_{\frac{1}{R_1^2}g}(p,1)$$

is foliated by $U(1)$-orbits (circles) and for every one, one can consider the area minimizing disc according to Proposition 2 and with boundary the orbit. This construction can be done for every point p in $\{\mathcal{P}\}\setminus\mathcal{T}(\partial\Sigma,\epsilon_1/2)$ which gives us the set of all the discs we are looking for. Now, let S be a stable and axisymmetric minimal surface embedded in Σ. If $S\cap\mathcal{T}(\{\mathcal{P}\},\epsilon_2)\neq\emptyset$ then there is $p\in\{\mathcal{P}\}$ such that

$$S\cap B_{\frac{1}{R_1^2}g}(p,L)\neq\emptyset$$

But then by Proposition 2, $S\cap B_{\frac{1}{R_1^2}g}(p,L)$ must be one of the discs we defined before. Finally note that if an axisymmetric compact and boundary-less surface intersects the axes $\{\mathcal{P}\}$ then it must do twice and the surface must be an sphere.

Item 3. By standard minimal surfaces estimates [5] the Gaussian curvature $\mathcal{K}$ of any disc D as in *item 4* and with respect to the induced metric from $\frac{1}{R_1^2}g$ is uniformly bounded. Consider now polar coordinates in D as defined in Section 2.3. Then, by Gauss-Bonet we have

$$2\pi-\frac{d\,l(s)}{d\,s}=\int_0^s\mathcal{K}(\bar{s})l(\bar{s})d\bar{s}$$

From this, the estimate $|l(s)-2\pi s|\leq c_0(\rho_0,\mu_0,\mu_1)s^2$ easily follows. □

Proof of Proposition 6: Let i be 1 or 2. Let $\{S_m\}$ be a sequence of axisymmetric spheres in Σ_i. Denote by Ω_m the (closed) set enclosed by S_H and S_m. Assume that the sequence is such that $\Omega_m\subset\Omega_{m+1}$ and $\mathrm{dist}(S_H,S_m)\to\infty$. Let $\{\delta_m\}$ be a positive sequence in such a way that (for every m) δ_m is sufficiently small that we have g-Gaussian coordinates well defined inside $\mathcal{T}(S_m,\delta_m)\setminus\Omega_m$. Thus on $\mathcal{T}(S_m,\delta_m)\setminus\Omega_m$ we can write

$$g=dr^2+h_m,$$

where $r(p)=dist(p,S_m)>0$ and $h_m(\partial_r,-)=h_m(-,\partial_r)=0$. Define $\Omega_{\delta_m}=\Omega_m\cup\mathcal{T}(S_m,\delta_m)$. On Ω_{δ_m} we define the axisymmetric metric g^*_m

$$g^*_m=\begin{cases} g & \text{on}\quad \Omega_m,\\ dr^2+f(r)^2h_m & \text{on}\quad \mathcal{T}(S_m,\delta_m)\setminus\Omega_m\end{cases}\tag{36}$$

and where $f(r)$ is the scalar function

$$f(r)=1+e^{-\frac{1}{r}+\frac{1}{\delta_m+\epsilon_m-r}}$$

where ϵ_m is, for every m, small enough to make the boundary $\{r=\delta_m\}$ strictly convex. Observe that $g^*_m=g$ on Ω_m and that $g^*_m\geq g$ on Ω_{δ_m} (because $f\geq 1$).

We define $\underline{A}^{\mathcal{N}}_{i,m}(\tilde{p})$, in the same way as $\underline{A}^{\mathcal{N}}_{i}(\tilde{p})$, as the infimum of the g^*_m-areas of the axisymmetric discs $D^{\mathcal{N}}(C)$, $C=\Pi^{-1}(\tilde{p})$, inside Ω_{δ_m}.

Claim 1: $\underline{A}^{\mathcal{N}}_{i,m}(\tilde{p})$ is realized by either

D1'. *The g^*_m-area of a disc $D^{\mathcal{N}}(C)=\Pi^{-1}(\gamma^{\mathcal{N}}(\tilde{p}))\subset\Omega_{\delta_m}$, or by,*

D2'. *The g^*_m-area of a disc $D^{\mathcal{S}}(C)=\Pi^{-1}(\gamma^{\mathcal{S}}(\tilde{p}))\subset\Omega_{\delta_m}$ plus A_H.*

The proof of the *claim 1* is as follows. Let $\{D^{\mathcal{N}}_j(C)\}=\{\Pi^{-1}(\beta^{\mathcal{N}}_j(\tilde{p}))\}$ be a g^*_m-area minimizing sequence of discs in Ω_{δ_m}, that is, a sequence for which we have

$$\lim A_{g^*_m}(D^{\mathcal{N}}_j)(C))=\lim \text{length}_{\bar{\tilde{g}}^*_m}(\beta^{\mathcal{N}}(\tilde{p}))=\underline{A}^{\mathcal{N}}_{i,m}(\tilde{p})$$

(where as we defined $\bar{\tilde{g}}$ we define here $\bar{\tilde{g}}^*_m:=\lambda^{*2}\tilde{g}^*_m$, $\lambda^{*2}=<\xi^*_m,\xi^*_m>_{g^*_m}$ the g^*_m-norm squared of the rotational Killing ξ^*_m of the metric g^*_m). On general grounds[9] the sequence of area minimizing discs converges in measure to a disc with boundary C (the solution of the "Plateau's problem") and, possibly, to a finite set of axisymmetric compact and non-contractible stable minimal surfaces, which because of the geometry of Ω_{δ_m} must be axisymmetric spheres. Thus the limit is either

P1. A stable minimal disc $D^{\mathcal{N}}(C)=\Pi^{-1}(\gamma^{\mathcal{N}}(\tilde{p}))$, or,

P2. A stable minimal disc $D^{\mathcal{S}}(C)=\Pi^{-1}(\gamma^{\mathcal{S}}(\tilde{p}))$, or,

P3. A stable minimal disc $D^{\mathcal{N}}(C)=\Pi^{-1}(\gamma^{\mathcal{N}}(\tilde{p}))$ plus a set of stable axisymmetric minimal spheres $S_k=\Pi^{-1}(\gamma^{\mathcal{N},\mathcal{S}}_k)$, $k=1,\ldots,k_1$, or,

P4. A stable minimal disc $D^{\mathcal{S}}(C)=\Pi^{-1}(\gamma^{\mathcal{S}}(\tilde{p}))$ plus a set of stable axisymmetric minimal spheres $S_k=\Pi^{-1}(\gamma^{\mathcal{N},\mathcal{S}}_k)$, $k=1,\ldots,k_2$.

Of course to guarantee the existence of the limit one uses that $(\Omega_{\delta_m},g^*_m)$ has one totally geodesic boundary (S_H) and one strictly convex boundary. We show now that the cases **P2** and **P3** cannot occur and that case **P3** could occur but when does then it does only with $k_2=1$ and $A_{g^*_m}(S_1)=\text{length}_{\bar{\tilde{g}}^*_m}(\gamma^{\mathcal{N},\mathcal{S}}_1(\tilde{p}))=A_H$. We show this below, which completes the proof of the *claim 1*.

Case **P3** cannot occur for in that case the (constant) sequence of curves $\{\beta'^{\mathcal{N}}_j(\tilde{p})=\gamma^{\mathcal{N}}(\tilde{p})\}$ has

$$\lim \text{length}_{\bar{\tilde{g}}^*_m}(\beta'^{\mathcal{N}}_j(\tilde{p})))<\lim \text{length}_{\bar{\tilde{g}}^*_m}(\beta^{\mathcal{N}}_j(\tilde{p}))$$

which is not possible because $\{\beta^{\mathcal{N}}_j(\tilde{p})\}$ is by assumption a minimizing sequence.

We show now that the case **P2** cannot occur. Let $\epsilon>0$ be small enough such that the curve $\gamma^{\mathcal{N}}(\tilde{p})$ does not intersects $\mathcal{T}_{\bar{\tilde{g}}^*_m}(\{\mathcal{N}\},\epsilon)$ where the tubular neighborhood is inside Ω^*_m. If the sequence $\{\beta^{\mathcal{N}}_j(\tilde{p})\}$ converges in measure to $\gamma^{\mathcal{N}}(\tilde{p})$ then it must be

$$\lim \text{length}_{\bar{\tilde{g}}^*_m}(\beta^{\mathcal{N}}_j(\tilde{p})\cap\mathcal{T}(\{\mathcal{N}\},\epsilon))=0$$

---

[9] Despite it naturalness this does not follow exactly from the well known result in Riemannian geometry that a minimizing sequence of curves in a complete boundaryless Riemannian manifold with fixed extreme points converges in measure to a geodesic because on one side the manifold $\tilde{\Sigma}$ has boundary and on the other hand the metric $\bar{\tilde{g}}^*_m$ is singular at the axis $\{\mathcal{P}\}$ (the boundary of $\tilde{\Sigma}$. Although with more work a proof can be given along these lines, a rigorous proof follows from the standard results on geometric measure theory on area minimizing sequences of discs [9], [8].

But every curve from $\tilde{p}$ to $\{\mathcal{N}\}$ must intersect the tubular neighborhood in a curve (or a set of curves) of total length at least ϵ. This gives a contradiction.

We finally analyze case **P4**. First we show that $k_2 = 1$. Assume by contradiction that $k_2 \geq 2$. Then one can consider a sequence of curves $\{{\beta'}^{\mathcal{N}}_j(\tilde{p})\}$ that would allow us to show that the sequence $\{\beta^{\mathcal{N}}_j(\tilde{p})\}$ was not minimizing. The sequence $\{{\beta'}^{\mathcal{N}}_j(\tilde{p})\}$ is constructed as follows. Let $\{q^1_j\}$ and $\{q^2_j\}$ be sequences of points in $\gamma^{\mathcal{S}}(\tilde{p})$ and $\gamma^{\mathcal{N},\mathcal{S}}_1$ respectively converging to points in the south axis $\{\mathcal{S}\}$ (i.e. the (south) end points of the curves $\gamma^{\mathcal{N}}(\tilde{p})$ and $\gamma^{\mathcal{N},\mathcal{S}}_1$ respectively). Then ${\beta'}^{\mathcal{N}}_j(\tilde{p})$ is defined (starting from $\tilde{p}$) as $\gamma^{\mathcal{N}}(\tilde{p})$ until q^1_j, then as a curve joining q^1_j and q^2_j going very near the south axis $\{\mathcal{S}\}$ and then as the piece of $\gamma^{\mathcal{N},\mathcal{S}}_{k_1}$ that starts at q^2_j and end at the north point of it. The curve between q^1_j and q^2_j has to be chosen in such a way that its $\bar{\tilde{g}}^*_m$-length goes to zero as $j \to \infty$ (note that the pre-image of such curve under Π becomes a very thin tube joining $\Pi^{-1}(q^1_j)$ and $\Pi^{-1}(q^2_j)$ of small g^*_m-area). With this definition of $\{{\beta'}^{\mathcal{N}}_j(\tilde{p})\}$ we have

$$\lim \mathrm{length}_{\bar{\tilde{g}}^*_m}({\beta'}^{\mathcal{N}}_j(\tilde{p})) = \mathrm{length}_{\bar{\tilde{g}}^*_m}(\gamma^{\mathcal{S}}(\tilde{p})) + \mathrm{length}_{\bar{\tilde{g}}^*_m}(\gamma^{\mathcal{N},\mathcal{S}}_1) < \lim \mathrm{length}_{\bar{\tilde{g}}^*_m}(\beta^{\mathcal{N}}_j(\tilde{p}))$$

which is impossible because by assumption $\{\beta^{\mathcal{N}}_j(\tilde{p})\}$ was a minimizing sequence. That $\mathrm{length}_{\bar{\tilde{g}}^*_m}(\gamma^{\mathcal{N},\mathcal{S}}_1) = A_H$ is a consequence of the fact that one can easily construct a sequence of curves $\{{\beta'}^{\mathcal{N}}(\tilde{p})\}$ (following a similar procedure as before) converging in measure to $\gamma^{\mathcal{S}}(\tilde{p}) \cup \gamma^{\mathcal{N},\mathcal{S}}_H$.

Claim2: For every $\tilde{p} \in \tilde{\Sigma}^\circ_i$ and constant $B > 0$ there is $m(B)$ such that for any $m \geq m(B)$ and g^*_m-stable axisymmetric minimal disc $D_m = D^{\mathcal{N}}_m(\Pi^{-1}(\tilde{p})) = \Pi^{-1}(\gamma^{\mathcal{N}}_m(\tilde{p}))$ or disc $D_m = D^{\mathcal{S}}_m(\Pi^{-1}(\tilde{p})) = \Pi^{-1}(\gamma^{\mathcal{S}}_m(\tilde{p}))$, inside Ω_{δ_m} and intersecting $S_{m(B)}$ we have

$$A_g(D_m \cap \Omega_{m(B)}) = A_{g^*_m}(D_m \cap \Omega_{m(B)}) \geq B$$

The Proof of the *claim 2* is as follows. Let m and $m(B)$ with $m \geq m(B)$ be arbitrary and let D_m be a disc as in the hypothesis. By Corollary 3, there is $\epsilon_2 > 0$ such that if D_m intersects $\mathcal{T}(\{\mathcal{P}\}, \epsilon_2) \setminus \mathcal{T}(S_{m(B)}, 1)$ then it intersects $\mathcal{T}(\{\mathcal{P}\}, \epsilon_2) \cap \Omega_{m(B)}$ exactly in a small disc. Therefore if $D_m \cap S_{m(B)} \neq \emptyset$ then $\gamma^{\mathcal{N}}_m(\tilde{p})$ or $\gamma^{\mathcal{S}}_m(\tilde{p})$ (depending on the case) remains inside $\big(\tilde{\Sigma}_i \setminus \Pi(\mathcal{T}(\{\mathcal{P}\}, \epsilon_2))\big) \cap \Pi(\Omega_{m(B)})$ until entering $\Pi(\mathcal{T}(S_{m(B)}, 1))$ for the first time. By the homogeneous regularity of the metric $\bar{\tilde{g}}$ on $\tilde{\Sigma} \setminus \Pi(\mathcal{T}(\{\mathcal{P}\}, \epsilon_2)$ we conclude that if $m(B) \to \infty$ then necessarily $\mathrm{length}_{\bar{\tilde{g}}}(\gamma^{\mathcal{N}}_m(\tilde{p}) \cap \Pi(\Omega_{m(B)})) \to \infty$. The claim follows then from the identity $A(\Pi^{-1}(\gamma^{\mathcal{N}}_m(\tilde{p})) \cap \Omega_{m(B)}) = \mathrm{length}_{\bar{\tilde{g}}}(\gamma^{\mathcal{N}}_m(\tilde{p}) \cap \Pi(\Omega_{m(B)}))$.

Now, let $\bar{D}^{\mathcal{N}}(C)$, $\Pi(C) = \tilde{p}$, be a fixed disc in Σ_i. Let $B = 2A(\bar{D}^{\mathcal{N}}(C))$ and $m(B)$ as in the *claim* (assume that $m(B)$ is big enough that $\bar{D}^{\mathcal{N}}(C) \subset \Omega_{m(B)}$). Then for any $m \geq m(B)$ we have

$$\underline{A}^{\mathcal{N}}_{i,m}(\tilde{p}) \leq A(\bar{D}^{\mathcal{N}}(C)) = \frac{B}{2}$$

It follows from the *claim 2* that for any $m \geq m(B)$ the minimizers $D^{\mathcal{N}}(C)$ (in case **D1'**) or $D^{\mathcal{S}}(C)$ (in case **D2'**) realizing $\underline{A}^{\mathcal{N}}_{i,m}(\tilde{p})$, lie in $\Omega_{m(B)}$. Therefore for any $m \geq m(B)$ we have

$$\underline{A}^{\mathcal{N}}_{i,m}(\tilde{p}) = \underline{A}^{\mathcal{N}}_{i,m(B)}(\tilde{p})$$

As $\lim_{m\to\infty} \underline{A}^{\mathcal{N}}_{i,m}(\tilde{p}) = \underline{A}^{\mathcal{N}}_{i}(\tilde{p})$ we conclude that the minimizers $D^{\mathcal{N}}(C)$ (in case **D1'**) or $D^{\mathcal{S}}(C)$ (in case **D2'**) realizing $\underline{A}^{\mathcal{N}}_{i,m}(\tilde{p})$ are the minimizers claimed **D1** or **D2** in the statement of the Proposition. The rest of the claim in the Proposition are automatic. □

References

[1] A. Aceña, S. Dain, and M. E. Gabach Clément. Horizon area-angular momentum inequality for a class of axially symmetric black holes. *Classical and Quantum Gravity*, 28(10):105014, May 2011.

[2] A. J. Amsel, G. T. Horowitz, D. Marolf, and M. M. Roberts. No dynamics in the extremal Kerr throat. *Journal of High Energy Physics*, 9:44, September 2009.

[3] Lars Andersson, Michael Eichmair, and Jan Metzger. Jang's equation and its applications to marginally trapped surfaces. In *Complex analysis and dynamical systems IV. Part 2*, volume 554 of *Contemp. Math.*, pages 13–45. Amer. Math. Soc., Providence, RI, 2011.

[4] N. Aronszajn. A unique continuation theorem for solutions of elliptic partial differential equations or inequalities of second order. *J. Math. Pures Appl. (9)*, 36:235–249, 1957.

[5] Tobias Holck Colding and William P. Minicozzi, II. *A course in minimal surfaces*, volume 121 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2011.

[6] S. Dain and M. Reiris. Area-Angular-Momentum Inequality for Axisymmetric Black Holes. *Physical Review Letters*, 107(5):051101, July 2011.

[7] J. L. Jaramillo, M. Reiris, and S. Dain. Black hole area-angular-momentum inequality in nonvacuum spacetimes. *Physical Review Letters D*, 84(12):121503, December 2011.

[8] William Meeks, III, Leon Simon, and Shing Tung Yau. Embedded minimal surfaces, exotic spheres, and manifolds with positive Ricci curvature. *Ann. of Math. (2)*, 116(3):621–659, 1982.

[9] Leon Simon. *Lectures on geometric measure theory*, volume 3 of *Proceedings of the Centre for Mathematical Analysis, Australian National University*. Australian National University Centre for Mathematical Analysis, Canberra, 1983.

[10] Robert M. Wald. *General relativity*. University of Chicago Press, Chicago, IL, 1984.